



Project group

Functional molecules on silicon surfaces

TPs involved (DEC'06):

B 6 (Rück-Braun)


B 1 (Tegeuder/Wolf)

A 4 (Raschke/Elsässer)

A 2 (Fölsch/Koch)

Si(111)




Self-assembled monolayers (SAM)



- B 6** (Design & synthesis)
- B 1** (Optical spectroscopy)
- A 4** (Near-field microscopy)

Si(100)

Single
1,5-cyclooctadiene
molecules



A 2 (LT-STM/STS)

Cluster calculations in TP **C 2**
(Saalfrank, Klamroth) in prep.

Self-assembled monolayers (SAM) on Si(111)

Characterization – current status:





- Surface tension measurement (**B 6**)
- AFM (in air), IR absorption spectroscopy (**A 4**)

Consensus: Further characterization and more analytical tools needed!

- Improvement of sensitivity in IR absorption (**A 4**)
- XPS and Sum-Frequency-Generation (SFG) (**B 1**)
- Photoelectron spectroscopy at BESSY (agreement with **B 2**, Weinelt/Gießel)
- LT-STM at LN temperature (**A 2**) (crucial points: surface contamination, sufficient conductance)

What we would wish to have:

Room-temperature STM to enable faster sample throughput



AFM of (C₁₀H₂₀CO₂Et SAM)
M. Breusing (**A 4**)