

## Statistical Physics and Thermodynamics (SS 2018)

### Problem sheet 3

**Hand in: Friday, May 18 during the lecture (note: one week later than usual)**

<http://www.physik.fu-berlin.de/en/einrichtungen/ag/ag-netz/lehre>

#### 1 Total differential (7 points)

The differential  $\alpha(x, y) = A(x, y)dx + B(x, y)dy$  is called *exact* if there is a function  $F(x, y)$ , so that  $dF(x, y) = (\partial F/\partial x)dx + (\partial F/\partial y)dy$  is equal to  $\alpha(x, y)$ .

a) Show that if  $\alpha$  is exact, then  $A(x, y)$  and  $B(x, y)$  satisfy the condition

$$\frac{\partial A(x, y)}{\partial y} = \frac{\partial B(x, y)}{\partial x}. \quad \text{(2 points)} \quad (1)$$

*Remark: The converse not necessarily holds.*

Consider  $\alpha(x, y) := (x^2 - xy)dx + x^2dy$

b) Does  $\alpha(x, y)$  fulfill equation (1)? **(1 point)**

c) Determine the exponent  $n \in \mathbb{N}$  so that  $\beta(x, y) = \alpha(x, y)/x^n$  is an exact differential. **(2 points)**

d) Find a function  $F(x, y)$  such that  $dF(x, y) = \beta(x, y)$  and  $F(1, 0) = 0$ . **(1 point)**

e) Find a curve  $(x, y(x))$  in space where  $F(x, y) = F(1, 0)$  is constant. Sketch  $y(x)$  in the range  $0 < x < 3$ . **(1 point)**

#### 2 Legendre transformation (3 points)

Consider a function  $f(x)$  and the differential  $df := u(x)dx$  where  $u(x) = df(x)/dx$ .

a) Show that you can define a function  $g(u) := xu(x) - f(x)$  for which,

$$\frac{dg}{du} = x \quad \text{(1 point)} \quad (2)$$

$g(u)$  is the Legendre transformed function of  $f(x)$ . We can also define the Legendre transform for a function of two variables  $g(x, v) := yv(x, y) - f(x, y)$ , where  $v(x, y) = \partial f(x, y)/\partial y$ .

b) Consider the two functions  $f_1(x) = \alpha x^2$  and  $f_2(x, y) = \alpha x^2 y^3$  where  $\alpha \in \mathbb{R}$  and calculate their Legendre transformations  $g_1(u)$  and  $g_2(x, v)$ . **(2 points)**

### 3 Phase space and 1D harmonic oscillator (10 points)

Consider a 1D harmonic oscillator with mass  $m$  and position  $q$  in a potential  $V(q) = kq^2/2$ .

- Write down the Lagrange function  $L(q, \dot{q}) = T - V$ , where  $T$  is the kinetic energy and  $V$  the potential energy, and derive the equation of motion for  $q$  using the Euler-Lagrange equation. **(1 point)**
- For the 1D harmonic oscillator, explicitly calculate the Legendre transformation

$$\mathcal{H}(q, p) = p\dot{q}(q, p) - L(q, \dot{q}(q, p)), \quad (3)$$

where the canonical momentum is defined by  $p = \partial L / \partial \dot{q}$ . **(1 point)**

- Calculate Hamilton's equations of motion, and solve them to obtain the phase space trajectory  $(q(t), p(t))$  with initial conditions  $q(0) = q_0$ ,  $p(0) = p_0$ . **(2 points)**
- Use your result from c) to calculate  $\mathcal{H}(q(t), p(t))$  and  $L(q(t), \dot{q}(t))$  along a solution of the equations of motion. Are they conserved along a trajectory? **(2 points)**
- Sketch a solution from c) with initial energy  $E = \mathcal{H}(q_0, p_0)$  in phase space, i.e. in the  $(q, p)$ -plane. Is the resulting curve closed? What geometric shape does the trajectory have? At which points does it intersect the  $q$ - and  $p$ -axes? **(2 points)**
- Assume the oscillator starts at rest with energy  $E$ . Calculate the running averages of kinetic and potential energy, i.e.

$$\bar{E}_{\text{kin}}(t) = \frac{1}{t} \int_0^t T(t') dt', \quad \bar{E}_{\text{pot}}(t) = \frac{1}{t} \int_0^t V(t') dt' \quad (4)$$

and express them in terms of  $E$ . What are the limits for  $t \rightarrow 0$ ,  $t \rightarrow \infty$ ? **(2 points)**