

Theory of Charge Sensing in Quantum-Dot Structures

Richard Berkovits,¹ Felix von Oppen,² and Yuval Gefen³

¹The Minerva Center, Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel

²Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany

³Department of Condensed Matter Physics, The Weizmann Institute of Science, Rehovot 76100, Israel

(Received 18 November 2004; published 23 February 2005)

Charge sensing in quantum-dot structures is studied by an exactly solvable reduced model and numerical density-matrix renormalization-group methods. Charge sensing is characterized by repeated cycling of the occupation of current-carrying states due to the capacitive coupling to trap states. In agreement with recent experiments, it results in characteristic asymmetric Coulomb-blockade peaks as well as sawtooth and domelike structures. Temperature introduces asymmetric smearing of these features and correlations in the conductance provide a fingerprint of charge sensing.

DOI: 10.1103/PhysRevLett.94.076802

PACS numbers: 73.23.Hk, 71.15.Dx, 73.23.-b

Introduction.—Within the orthodox picture of the Coulomb blockade, subsequent Coulomb-blockade peaks are due to the filling of consecutive single-particle states [1]. Once a state is filled, it remains so. Although this picture successfully describes various transport properties of weakly coupled nanoscopic systems, there has been much interest in identifying situations in which the orthodox picture fails and a “dynamical” behavior of the occupations of single-particle orbitals emerges.

One of the earliest pertinent examples was pointed out by Kuznetsov *et al.* [2] who considered the filling of localized states in a barrier. They have shown that as the gate voltage increases, a localized state may first fill and then vacate, once a different localized state is occupied. This behavior is manifested in the conduction through the barrier by the reappearance of the *same* conduction peak. A new wave of interest was motivated by the correlations observed in the transmission phase through a quantum dot [3]. An attractive explanation for these correlations is that a number of successive transmission peaks through the dot is carried by the same state [4–6]. This, of course, requires the population of this state to be repeatedly cycled. Various mechanisms, which lie beyond the orthodox picture, have been proposed [4–6].

In this paper, we show that repeated filling of a single-particle state is in fact a rather generic phenomenon. We find that it occurs whenever there exist traps in the system, either by accident or by specific design of a quantum-dot structure. This phenomenon has been seen by Lindemann *et al.* [7] and in tailored structures by Johnson *et al.* [8] and Kobayashi *et al.* [9]. Consistent with these experiments, we observe that repeated filling of a given single-particle state as a function of gate voltage can be reflected in the conductance in many different ways, ranging from essentially no signature to sawtooth or domelike structures to asymmetric Coulomb-blockade peaks. The underlying mechanism termed *charge sensing* in Ref. [8], is based on the capacitive coupling between the traps and the conducting channel.

We first obtain our results within a reduced, exactly solvable model which captures the essential features of the phenomenon. As a by-product we show how the mechanism previously proposed by Silvestrov and Imry [6] is a limiting case of our more general approach. Relying on numerical results obtained by a density-matrix renormalization-group (DMRG) method [10,11], we subsequently discuss how our results are modified when relaxing various restrictions of the exactly solvable model.

Model.—Motivated by the experiment of Ref. [8], we consider a reduced model of the charge-sensing setup as shown in Fig. 1. A quantum dot is coupled to two leads. This *connected* quantum dot is coupled electrostatically to a *disconnected* dot in its vicinity. Within our reduced model, we make the following assumptions: (i) There is no tunneling between the disconnected dot and the leads or




FIG. 1. Conductance (solid line) through one connected state with ten disconnected states as function of chemical potential. (a) Intermediate coupling case: $\epsilon = -1.08$, $t = 0.2$, $U = 0.16$, and $e_m = -1.2 + 0.02m$ ($m = 1, \dots, 10$). (b) Strongly connected case: $\epsilon = -1$, $t = 0.5$, $U = 0.08$, $e_m = -1.24 + 0.04m$. Dotted curves: Conductances for fixed occupations $n_d = 0, \dots, 10$ of the disconnected states. As n_d changes with μ , the conductance (solid line) switches accordingly. Inset: Schematic of the connected and disconnected dots.

the connected dot. (ii) Transport through the connected dot is carried by a single state. (iii) The disconnected dot may have many single-particle levels. We emphasize that this model may also represent a quantum point contact with nearby traps, as well as transport through a single quantum dot in which one state is much more strongly coupled to the leads than the others [6,7]. In the latter case, transport through the weakly coupled states does not contribute significantly to the conductance. Instead, the weakly coupled states predominantly affect the system through their capacitive coupling to the well coupled state and may thus, to leading order, be thought of as trap states.

Our reduced model is defined by the Hamiltonian

$$H = H_{\text{dots}} + H_{\text{leads}} + H_{\text{mix}}. \quad (1)$$

Here, the Hamiltonian H_{dots} of the dots involves the level energy ϵ of the connected dot (with creation operator α^\dagger) and e_m of the disconnected dot (with creation operator d_m^\dagger) together with the charging energies U_d and U_m for the disconnected dot and for the mutual capacitive coupling of both dots. For simplicity, we will assume these charging energies to be equal. For spinless electrons this gives $H_{\text{dots}} = \epsilon \alpha^\dagger \alpha + \sum_{m=1}^{N_d} e_m d_m^\dagger d_m + U[n_d(n_d - 1)/2 + n_\alpha n_d]$, where $n_\alpha = \alpha^\dagger \alpha$ and $n_d = \sum_{m=1}^{N_d} d_m^\dagger d_m$. The lead Hamiltonian is $H_{\text{leads}} = \sum_{k,\lambda=L,R} E_{k\lambda} c_{k\lambda}^\dagger c_{k\lambda}$ and the tunneling between lead and connected dot is described by $H_{\text{mix}} = \sum_{k\lambda} t_{k\lambda} \alpha^\dagger c_{k\lambda} + \text{H.c.}$

Generally, Eq. (1) is a many-particle Hamiltonian. Note, however, that there is no charging term for the connected dot since it can be only singly occupied. When combined with the assumption of no tunneling to and from the disconnected dot, this fact makes our reduced model *exactly solvable*. Since the occupations $\langle d_m^\dagger d_m \rangle$ of the disconnected-dot states can only take on the values 0 or 1, we can treat these operators as c numbers in the Hamiltonian Eq. (1). Thus, the Hamiltonian can be regarded as a set of 2^{N_d} *single-particle* Hamiltonians, one for each possible set $\{n_m\}$ of occupation numbers $d_m^\dagger d_m$,

$$H_{\{n_m\}} = (\epsilon + U n_d) \alpha^\dagger \alpha + \sum_m e_m n_m + U n_d (n_d - 1)/2 + H_{\text{leads}} + H_{\text{mix}}. \quad (2)$$

The corresponding thermodynamic potentials at a given chemical potential μ and zero temperature are

$$\Omega_{\{n_m\}} = H_{\{n_m\}} - \mu(\alpha^\dagger \alpha + n_d + \sum_{k,\lambda=L,R} c_{k\lambda}^\dagger c_{k\lambda}). \quad (3)$$

The ground state occupation $\{n_m\}$ of the dots can now be found by determining the configuration with the lowest thermodynamic potential. Since $\Omega_{\{n_m\}}$ is a single-particle thermodynamic potential, this can be done by calculating $\Omega_{\{n_m\}}$ for all 2^{N_d} possible configurations.

Analytical treatment.—Focusing on the essential physics, we present a full analytic treatment of the thermodynamic potentials in Eq. (3) for a *single* disconnected state of energy e_1 . The extension to several disconnected states

is straightforward. For a single disconnected state, there are two different possibilities $n_d = 0$ and $n_d = 1$ with corresponding thermodynamic potentials Ω_0 and Ω_1 . As the chemical potential (gate voltage) μ increases, the disconnected state will be filled at $\mu = \mu_{\text{switch}}$ when the condition $\Omega_0 = \Omega_1$ is fulfilled. This switch in the occupation of the disconnected state is accompanied by an abrupt change in the occupation of—and hence the conductance through—the connected dot. It is this general phenomenon that is referred to as *charge sensing*. We now proceed with a quantitative analysis.

The Hamiltonians H_0 and H_1 , associated with empty and occupied disconnected state, respectively, are single-particle Hamiltonians with eigenstates $|\psi_j^0\rangle$ and $|\psi_j^1\rangle$ and eigenenergies $\{\epsilon_j^0\}$ and $\{\epsilon_j^1\}$. Thus, we can define the density of states on the connected dot $\nu^{0(1)}(\epsilon) = \sum_j \langle \psi_j^{0(1)} | \alpha^\dagger \alpha | \psi_j^{0(1)} \rangle \delta(\epsilon - \epsilon_j^{0(1)})$, and on the leads $\mathcal{N}^{0(1)}(\epsilon) = \sum_j \langle \psi_j^{0(1)} | \sum_{k\lambda} c_{k\lambda}^\dagger c_{k\lambda} | \psi_j^{0(1)} \rangle \delta(\epsilon - \epsilon_j^{0(1)})$. We now express the thermodynamic potentials as

$$\Omega_{n_d} = (e_1 - \mu) \delta_{n_d,1} + \int_{-\infty}^{\mu} d\epsilon (\epsilon - \mu) (\nu^{n_d}(\epsilon) + \mathcal{N}^{n_d}(\epsilon)).$$

Assuming that the density of states of the lead varies on scales large compared to both U and the width Γ of the connected state, we can write $\nu^0(\epsilon) = (\Gamma/2\pi)((\epsilon - \epsilon)^2 + (\Gamma/2)^2)^{-1}$ and $\nu^1(\epsilon) = \nu^0(\epsilon - U)$. Moreover, a finite set of disconnected states leaves the continuum of lead states essentially unaffected so that $\mathcal{N}^0 = \mathcal{N}^1$. Thus, we obtain

$$\begin{aligned} \Omega_1 - \Omega_0 = & e_1 - \mu + U/2 + \frac{\mu - \epsilon}{\pi} \arctan\left(\frac{2(\mu - \epsilon)}{\Gamma}\right) \\ & - \frac{\mu - U - \epsilon}{\pi} \arctan\left(\frac{2(\mu - U - \epsilon)}{\Gamma}\right) \\ & + \frac{\Gamma}{4\pi} \ln\left(\frac{(\mu - U - \epsilon)^2 + (\Gamma/2)^2}{(\mu - \epsilon)^2 + (\Gamma/2)^2}\right) \end{aligned} \quad (4)$$

upon performing the integration.

We first consider the limit of a *weakly connected* dot for which the distances of the connected-dot energies from the chemical potential $\epsilon - \mu$ (for $n_d = 0$), and $\epsilon + U - \mu$ (for $n_d = 1$) are large compared to the level width Γ . Then, Eq. (4) simplifies to $\Omega_1 - \Omega_0 \simeq e_1 - \epsilon + (\Gamma/2\pi) \times \ln(|\mu - \epsilon - U|/|\mu - \epsilon|)$, in agreement with the many-body perturbation theory result of Silvestrov and Imry [6]. This implies that the switching occurs on the Coulomb-blockade plateau ($\langle n_\alpha \rangle$ is an integer); hence there is no effect on the conductance at μ_{switch} . However, the switch leads to repeated appearances of the same Coulomb-blockade peak [6].

We now turn to the situation when the broadening Γ is comparable (*intermediate* coupling) or larger (*strongly connected*) than the distances of the dot energies from the chemical potential μ . In these cases, the perturbation theory of Ref. [6] fails, while our general solution Eq. (4) still applies. Specifically as μ sweeps across μ_{switch} from

below, the occupation of the connected-dot state decreases abruptly from $n_\alpha = 1/2 + \arctan[2(\mu_{\text{switch}} - \epsilon)/\Gamma]/\pi$ to $n_\alpha = 1/2 + \arctan[2(\mu_{\text{switch}} - \epsilon - U)/\Gamma]/\pi$. In the limit of a strongly connected state, Eq. (4) leads to the explicit solution $\mu_{\text{switch}} \simeq e_1 + U/2$.

Instead of the occupations n_α , we focus on the experimentally accessible (dimensionless) conductances g of the quantum dot. For the case of a quantum dot with two symmetrically coupled single-channel leads, the Friedel sum rule implies the relation $g = \sin^2(\pi n_\alpha)$ in the usual way [12]. Thus, the jump in occupation n_α at μ_{switch} translates directly into a jump in conductance (unless $\Delta n_\alpha = 1$, as happens for a weakly connected dot).

Numerical results.—Representative traces of the conductance as a function of μ are shown in Fig. 1. These plots are based on a generalization of the above analytic results to the case of an arbitrary number N_d of disconnected states. The most striking behavior occurs for intermediate coupling where the broadening Γ is comparable to or slightly smaller than the charging energy U . In this case shown in Fig. 1(a), one observes the appearance of *new asymmetric* peaks in the conductance trace. In the absence of disconnected states, there would be only a single conductance peak due to the single level of the connected dot. In the presence of the disconnected states, the occupation of the connected state decreases abruptly whenever a disconnected state is filled up. Thus the sharp jump in the conductance is downward (upward) if it occurs on the rising (falling) side of the conductance peak of the connected level. This leads to the appearance of new trap-induced peaks in g whose asymmetry arises from the abrupt jumps. Indeed, hints of this behavior have recently been seen in experiments and were attributed to charging of disconnected states [7].

For strongly connected dots ($U \ll \Gamma$), the jumps in the occupation n_α of the connected states which are associated with charging of disconnected states are typically small compared to one. In this case, also the abrupt changes in the conductance are small compared to the conductance itself, leading to a characteristic sawtooth-behavior of g as function of chemical potential. This is shown in Fig. 1(b). When the connected level is close to half filling, a typical dome shape is observed. This is very similar to the behavior seen in recent charge-sensing experiments on a quantum point contact monitoring the charge in a disconnected dot [8].

In experiments, a gate voltage does not affect connected and disconnected levels in the same way. Using different gates one can even manipulate the levels independently [8]. Such experiments correspond to nonvertical trajectories in the (V_g, μ) plane where μ is the overall chemical potential and V_g is assumed to affect the disconnected states only. The resulting intricate pattern of the conductance in the (V_g, μ) plane is shown in Fig. 2.

Temperature leads to very interesting behavior of the conductance, even when $kT \ll \Gamma$. In the latter regime, the sharp jumps of the conductance are broadened by tempera-




FIG. 2 (color online). Conductance as a function of overall chemical potential μ and external gate voltage V_g for the disconnected states (see inset in Fig. 1). Connected state: $\epsilon = 0$; disconnected states: $e_m = -0.55 + 0.05m$ ($m = 1, \dots, 5$), $U = 0.1$, and $t = 0.2$.

ture, while the other side of the conductance peaks is smooth on the scale Γ and thus insensitive to T . This results in a very asymmetric temperature broadening of the sawtooth peaks which was indeed observed in experiments [7]. These considerations can be quantified by noting that close to the charging of a disconnected state the system can be well approximated by a two-level system, with the two levels corresponding to $n_d = 0$ and $n_d = 1$ (or, more generally, $n_d = N$ and $n_d = N + 1$). The occupation of the connected state is then given by

$$n_\alpha = \frac{[\frac{1}{2} + \arctan\frac{2\mu}{\Gamma}]e^{-\omega_0} + [\frac{1}{2} + \arctan\frac{2(\mu-U)}{\Gamma}]e^{-\omega_1}}{e^{-\omega_0} + e^{-\omega_1}},$$

where $\omega_{n_d} = \Omega_{n_d}/kT$. Since the entropy is governed by the lead states which are unaffected by the change in n_d , one expects the entropy terms in Ω to be equal, i.e., $TS_1 = TS_0$, and thus to cancel out. Expanding $\Omega_1(\mu_{\text{switch}} + \delta\mu) - \Omega_0(\mu_{\text{switch}} + \delta\mu) \sim kT$ in $\delta\mu$, one obtains $kT \sim \delta\mu \{1 - \pi^{-1}[\arctan(2(\mu_{\text{switch}} - U)/\Gamma) - \arctan(2\mu_{\text{switch}}/\Gamma)]\}$. Therefore, the abrupt change in n_α and g is smeared by temperature over a range kT in μ . The resulting T dependence of g is depicted in Fig. 3(a).

A qualitatively similar effect occurs when the disconnected state is broadened, e.g., by coupling it to an external reservoir different from the current-carrying leads. This situation is no longer amenable to an exact solution since the occupation of the “disconnected” state can now differ from 0 or 1, and we resort to a numerical DMRG method. [We have checked that the DMRG reproduces our exact solution for vanishing broadening; see Fig. 3(b).] We find that the main effect of a finitely coupled disconnected state is to smear the sudden jump in the occupation of the latter, similar to the effect of finite temperature. Correspondingly, the effect of the broadening on g is very similar to that of temperature, as can indeed be verified by comparing Figs. 3(a) and 3(c).

This observation enables us to qualitatively understand the physics of two connected states of the same dot with




FIG. 3. (a) Temperature dependence of the conductance as function of μ (connected state: $\epsilon = -1.3$; disconnected state: $e_1 = -1.26$; $U = 0.4$; $t = 0.3$). The different curves correspond to $\pi kT = 0, 0.1, 0.2, 0.3, 0.4$. (b) Occupation of the connected state n_α (open symbols) and disconnected state n_d (filled symbols) for several values of couplings as the disconnected state opens up *additional* non-current-carrying lead. Lines: results of the exact solution of Eq. (3); symbols: DMRG calculations for different couplings of the disconnected state (circles $t = 0$, squares $t = 0.1$, diamonds $t = 0.2$). (c) Conductance of connected state in (b).

$\Gamma_1 \gg \Gamma_2$. Indeed, the level occupations are insensitive to whether the levels are coupled to the same or different leads. Figure 3(b) then applies and is consistent with very recent works on such setups employing Hartree-Fock and numerical RG methods [13,14]. Evidently, the way these states are coupled to the leads is crucial for the conductance, due to Fano-type interference effects [8,9].

The insight gained from our analysis involving a single connected state may be used to make quantitative predictions concerning *several* connected and disconnected states, including charging energies for the connected-dot states. Although it is then difficult to calculate the exact switching point for a particular disconnected state, one nevertheless predicts that for chemical potentials μ (or V_g) immediately before or after the switching, $n_{\alpha j}(\mu \pm 0) = n_{\alpha j}(\mu \mp U)$ and therefore $g(\mu \pm 0) = g(\mu \mp U)$. (Here, $n_{\alpha j}$ denotes the occupation of the j th connected state.) This is illustrated in Fig. 4 for two connected and two disconnected states, based on a DMRG calculation. This prediction should be very useful for analyzing experimental data. For any abrupt jump due to charging of a disconnected state, the conductance satisfies this relation. By contrast, if the jump in the conductance is due, e.g., to noise, no such correlation is expected.

In conclusion, it is interesting to speculate that the charge-sensing physics discussed theoretically in this Letter may occur generically in relatively well coupled chaotic quantum dots. By the nature of the Porter-Thomas distribution of lead-induced level broadenings, there will be a significant number of narrow levels in addition to broader levels, in particular, for systems with




FIG. 4. Occupation for two connected states ($t = 0.5$) at $\epsilon_1 = -1.1$ (gray circles) and $\epsilon_2 = -1$ (gray squares) and two disconnected states $e_1 = -1.12$ (solid line), $e_2 = -1.02$ (dotted line) as function of μ obtained from DMRG ($U = 0.3$). To demonstrate that right after the switch $n_\alpha(\mu) = n_\alpha(\mu - U)$, we plot $n_\alpha(\mu + U)$ for values of μ prior to the switch, i.e., $-1.2 < \mu < -1$ and $-0.48 < \mu < -0.28$ (white symbols). A clear overlap between the white and gray symbols is seen.

time-reversal symmetry. Charge sensing and patterns not unlike Fig. 1(b) may thus be important ingredients in explaining the large-scale structure of Coulomb-blockade sequences observed in such systems [15].

We acknowledge very useful discussions with I. Lerner, C. M. Marcus, Y. Oreg, and P. Silvestrov, as well as support from the Israel Science Foundation (Y. G. and R. B.), SFB 290, the JA (F.v.O.), and the BSF (Y.G.). One of us (F.v.O.) thanks the Einstein Center (Weizmann) and the Minerva Center (Bar-Ilan) for hospitality.

-
- [1] For a recent review, see Y. Alhassid, Rev. Mod. Phys. **72**, 895 (2000).
 - [2] V. V. Kuznetsov *et al.*, Phys. Rev. B **54**, 1502 (1996).
 - [3] A. Yacoby *et al.*, Phys. Rev. Lett. **74**, 4047 (1995); R. Schuster *et al.*, Nature (London) **385**, 417 (1997).
 - [4] Y. Oreg and Y. Gefen, Phys. Rev. B **55**, 13726 (1997).
 - [5] G. Hackenbroich, W.D. Heiss, and H. A. Weidenmüller, Phys. Rev. Lett. **79**, 127 (1997); R. Baltin *et al.*, Eur. Phys. J. B **10**, 119 (1999).
 - [6] P. Silvestrov and Y. Imry, Phys. Rev. Lett. **85**, 2565 (2000).
 - [7] S. Lindemann *et al.*, Phys. Rev. B **66**, 161312 (2002).
 - [8] A. C. Johnson *et al.*, Phys. Rev. Lett. **93**, 106803 (2004).
 - [9] K. Kobayashi *et al.*, Phys. Rev. B **70**, 035319 (2004).
 - [10] S. R. White, Phys. Rev. B **48**, 10345 (1993).
 - [11] R. Berkovits, F. von Oppen, and J.W. Kantelhardt, Europhys. Lett. **68**, 699 (2004).
 - [12] S. Datta and W. Tian, Phys. Rev. B **55**, 1914 (1997); M. Pustilnik and L.I. Glazman, Phys. Rev. Lett. **87**, 216601 (2001).
 - [13] M. Sindel *et al.*, cond-mat/0408096.
 - [14] J. König and Y. Gefen, cond-mat/0408691.
 - [15] S. M. Cronenwett *et al.*, Phys. Rev. Lett. **81**, 5904 (1998).