

The Nested Off-shell Bethe ansatz and $O(N)$ Matrix Difference Equations

Hrachya M. Babujian^{*†}, Angela Foerster^{‡§}, and Michael Karowski[¶]
 Institut für Theoretische Physik, Freie Universität Berlin,
 Arnimallee 14, 14195 Berlin, Germany

April 17, 2012

Abstract

A system of $O(N)$ -matrix difference equations is solved by means of the off-shell version of the nested algebraic Bethe ansatz. In the nesting process a new object, the Π -matrix, is introduced to overcome the complexities of the $O(N)$ group structure. The proof of the main theorem is presented in detail. In particular, the cancellation of all “unwanted terms” is shown explicitly. The highest weight property of the solutions is proved.

1 Introduction

$O(N)$ Gross-Neveu and $O(N)$ σ -models are asymptotically free quantum field theories which attract high interest, since they share some common features with QCD. Since perturbation theory fails for these models, exact results, such as exact generalized form factors are desirable and welcome. The concept of a generalized form factor was introduced in [1, 2], where several consistency equations were formulated. Subsequently this approach was developed further and investigated in different models by Smirnov [3]. Generalized form factors are matrix elements of fields with many particle states. To construct these objects explicitly one has to solve generalized Watson’s equations which are matrix difference equations. To solve these equations the so called “off-shell Bethe ansatz” is applied [4, 5, 6]. The conventional Bethe ansatz introduced by Bethe [7] is used to solve eigenvalue problems and its algebraic formulation was developed by Faddeev and coworkers (see e.g. [8]). The off-shell Bethe ansatz has been introduced in [9] to solve the

^{*}Address: Yerevan Physics Institute, Alikhanian Brothers 2, Yerevan, 375036 Armenia

[†]E-mail: babujian@yerphi.am

[‡]Address: Instituto de Física da UFRGS, Av. Bento Gonçalves 9500, Porto Alegre, RS - Brazil

[§]E-mail: angela@if.ufrgs.br

[¶]E-mail: karowski@physik.fu-berlin.de

Knizhnik-Zamolodchikov equations which are differential equations. In [10] a variant of this technique has been formulated to solve matrix difference equations of the form

$$K(u_1, \dots, u_i + \kappa, \dots, u_n) = K(u_1, \dots, u_i, \dots, u_n)Q(u_1, \dots, u_n; i), \quad (i = 1, \dots, n),$$

where $K(\underline{u})$ is a co-vector valued function, $Q(\underline{u}, i)$ are matrix valued functions and κ is a constant to be specified. We use here a co-vector formulation because this is more convenient for the application to the form factor program. For higher rank internal symmetry groups the nested version of this Bethe ansatz has to be applied. The nested Bethe ansatz as a method to solve eigenvalue problems was introduced by Yang [11] and further developed by Sutherland [12, 13].

In this article we will solve the $O(N)$ difference equations combining the nested Bethe ansatz with the off-shell Bethe ansatz. This procedure is similar to the $SU(N)$ [14] case, where also a nesting procedure is used. However, the algebraic formulation for $O(N)$ is much more intricate because the R-matrix exhibits an extra new term. In addition, for $SU(N)$ we can use the same R-matrix at every level, while for the group $O(N)$ the R-matrix changes after each level. Therefore in our construction a new object, called Π -matrix, is introduced in order to overcome these difficulties. This provides a systematic formulation of techniques introduced by Tarasov [15] and also used in [16]. In [17] a different procedure was used to solve the $O(N)$ on-shell Bethe ansatz for even N .

The results of this article will be applied in [18] to calculate exact form factors of the $O(N)$ σ - and Gross-Neveu models. We should mention that the first computation of form factors for $O(3)$ σ -model is due to [3] (see also [19, 20]). There are also new developments concerning the connection between 2d Conformal Field Theory (CFT) and integrable models with $N = 2$ Super Yang Mills (SYM) theories in different higher dimensions. First, there is a surprising relation between 2d-conformal blocks and the instanton partition function in $N = 2$, 4d-SYM theory [21] (Alday, Gaiotto, Tachikawa - AGT relation) and this is a particular version of the AdS/CFT correspondence which is a more general part of the gauge/string duality. There is also a q-deformation of the AGT relation which connects the $N = 2$ 5d-SYM theory and the q-deformed conformal blocks [22]. This last relation offers new insights and gives the intriguing hope that the form factor program can be used to obtain a deeper understanding of this connection. The solution of the difference equations is the first step to obtain the exact form factors and therefore important physical relations and correlation functions for integrable models. In fact, difference equations play a significant role in various contexts of mathematical physics (see e.g. [23] and references therein).

The article is organized as follows. In Section 2 we recall some results and fix the notation concerning the $O(N)$ R-matrix, the monodromy matrix and some commutation rules. We also introduce a new object, which we call the Π -matrix and which is a central element in our construction of the nested off-shell Bethe vector. In Section 3 we introduce the nested generalized Bethe ansatz to solve a system of $O(N)$ difference equations and present the solutions in terms of ‘‘Jackson-type Integrals’’. We introduce a new type of monodromy matrix fulfilling a new type of Yang-Baxter relation and which is adapted to the difference problem. In particular this yields a relatively simple proof of our main result, which is Theorem 3.5. In Section 4 we prove the highest weight property of the

solutions and calculate the weights. The appendices provide the more complicated proofs of the results we have obtained. In particular, in Appendix B we determine all “unwanted terms” in the Bethe ansatz and show that they cancel.

2 General setting and notion of the Π -matrix

2.1 The $O(N)$ - R-matrix

Let $V^{1\dots n}$ be the tensor product space

$$V^{1\dots n} = V_1 \otimes \dots \otimes V_n, \quad (1)$$

where the vector spaces $V_i \cong \mathbf{C}^N$, ($i = 1, \dots, n$) are copies of the fundamental vector representation space of $O(N)$ with the (real) basis vectors

$$|\alpha\rangle_r \in V_i, \quad (\alpha = 1, \dots, N).$$

It is straightforward to generalize the results of this paper to the case where the V_i are vector spaces for other representations. We denote the canonical basis vectors of $V^{1\dots n}$ by

$$|\alpha_1, \dots, \alpha_n\rangle \in V^{1\dots n}, \quad (\alpha_i = 1, \dots, N). \quad (2)$$

A vector $v^{1\dots n} \in V^{1\dots n}$ is given in terms of its components by

$$v^{1\dots n} = \sum_{\underline{\alpha}} |\alpha_1, \dots, \alpha_n\rangle_r v^{\alpha_1, \dots, \alpha_n}. \quad (3)$$

A matrix acting in $V^{1\dots n}$ is denoted by

$$A_{1\dots n} : V^{1\dots n} \rightarrow V^{1\dots n}. \quad (4)$$

We will also use the dual space $V_{1\dots n} = (V^{1\dots n})^\dagger$.

The $O(N)$ spectral parameter dependent R-matrix was found by Zamolodchikov-Zamolodchikov [24]¹. It acts on the tensor product of two (fundamental) representation spaces of $O(N)$. It may be written as

$$R_{12}(u_{12}) = (\mathbf{1}_{12} + c(u_{12}) \mathbf{P}_{12} + d(u_{12}) \mathbf{K}_{12}) : V^{12} \rightarrow V^{21}, \quad (5)$$

where \mathbf{P}_{12} is the permutation operator, \mathbf{K}_{12} the annihilation-creation operator and $u_{12} = u_1 - u_2$. Here and in the following we associate to each space V_i a variable (spectral parameter) $u_i \in \mathbb{C}$. The components of the R-matrix are

$$R_{\alpha\beta}^{\delta\gamma}(u_{12}) = \delta_\alpha^\gamma \delta_\beta^\delta + \delta_\alpha^\delta \delta_\beta^\gamma c(u_{12}) + \delta^{\gamma\delta} \delta_{\alpha\beta} d(u_{12}) = \begin{array}{ccc} & \delta & \gamma \\ & \diagdown & \diagup \\ u_1 & & u_2 \\ & \diagup & \diagdown \\ & \alpha & \beta \end{array}, \quad (6)$$

¹We use here the normalization $R = S/\sigma_2$ and the parameterization $u = \theta/i\pi\nu$ which is more convenient for our purpose.

from which \mathbf{P}_{12} and \mathbf{K}_{12} can be read off. The functions

$$c(u) = \frac{-1}{u}, \quad d(u) = \frac{1}{u - 1/\nu}, \quad \nu = \frac{2}{N - 2} \quad (7)$$

are obtained as the rational solution of the Yang-Baxter equation

$$R_{12}(u_{12}) R_{13}(u_{13}) R_{23}(u_{23}) = R_{23}(u_{23}) R_{13}(u_{13}) R_{12}(u_{12}) \quad (8)$$

$$\begin{array}{c} \diagup \quad \diagdown \\ | \quad | \\ \diagdown \quad \diagup \\ 1 \quad 2 \quad 3 \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ 1 \quad 2 \quad 3 \end{array}, \quad (9)$$

where we have employed the usual notation [11]. We will also use

$$\tilde{R}(u) = R(u)/a(u)$$

with

$$a(u) = 1 + c(u) = \frac{u - 1}{u}.$$

The ‘‘unitarity’’ of the R-matrix reads as

$$\tilde{R}_{21}(u_{21}) \tilde{R}_{12}(u_{12}) = 1 : \quad \begin{array}{c} \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ 1 \quad 2 \end{array} = \begin{array}{c} | \quad | \\ 1 \quad 2 \end{array}$$

and the three eigenvalues of the R-matrix are

$$R_{\pm}(u) = 1 \pm c(u) = \frac{u \mp 1}{u}, \quad R_0 = 1 + c(u) + Nd(u) = \frac{u + 1}{u} \frac{u + 1/\nu}{u - 1/\nu}. \quad (10)$$

The crossing relation may be written as

$$R_{12}(u_{12}) = \mathbf{C}^{2\bar{2}} R_{\bar{2}1}(\hat{u}_{12}) \mathbf{C}_{22} = \mathbf{C}^{1\bar{1}} R_{\bar{1}2}(\hat{u}_{12}) \mathbf{C}_{\bar{1}1} \quad (11)$$

$$\begin{array}{c} \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ 1 \quad 2 \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ | \quad | \\ \diagdown \quad \diagup \\ 1 \quad 2 \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ | \quad | \\ \diagup \quad \diagdown \\ 1 \quad 2 \end{array}, \quad (12)$$

where $\hat{u} = 1/\nu - u$. Here $\mathbf{C}^{1\bar{1}}$ and $\mathbf{C}_{\bar{1}1}$ are the charge conjugation matrices. Their matrix elements are $\mathbf{C}^{\alpha\bar{\beta}} = \mathbf{C}_{\alpha\bar{\beta}} = \delta_{\alpha\beta}$ where $\bar{\beta}$ denotes the anti-particle of β . In the real basis used up to now the particles are chargeless which means that $\bar{\beta} = \beta$ and \mathbf{C} is diagonal.

In the following we will use instead of the real basis $|\alpha\rangle_r$, ($\alpha = 1, 2, \dots, N$) the complex basis given by

$$\begin{aligned} |\alpha\rangle &= \frac{1}{\sqrt{2}} (|2\alpha - 1\rangle_r + i|2\alpha\rangle_r) \\ |\bar{\alpha}\rangle &= \frac{1}{\sqrt{2}} (|2\alpha - 1\rangle_r - i|2\alpha\rangle_r) \end{aligned}$$

for $\alpha = 1, 2, \dots, [N/2]$. If N is odd there is in addition $|0\rangle = |\bar{0}\rangle = |N\rangle_r$. The weight vector $w = (w_1, \dots, w_{[N/2]})$ and the charges of the one-particle states are given by

$$\begin{aligned} \text{for } |\alpha\rangle &: w_k = \delta_{k\alpha} \quad , \quad Q = 1 \\ \text{for } |\bar{\alpha}\rangle &: w_k = -\delta_{k\alpha} \quad , \quad Q = -1 \\ \text{for } |0\rangle &: w_k = 0 \quad , \quad Q = 0. \end{aligned}$$

Remark 2.1 For even N this means that we consider $O(N)$ as a subgroup of $U(N/2)$ and the charge Q is its $U(1)$ charge. For $N = 3$ we may identify the particles $1, \bar{1}, 0$ with the pions π_{\pm}, π_0 .

The highest weight eigenvalue of the R-matrix is

$$R_{11}^{11}(u) = R_+(u) = a(u).$$

We order the states as: $1, 2, \dots, (0), \dots, \bar{2}, \bar{1}$ (0 only for N odd). Then the charge conjugation matrix in the complex basis is of the form

$$\begin{aligned} \mathbf{C}^{\delta\gamma} &= \delta^{\delta\bar{\gamma}}, \quad \mathbf{C}_{\alpha\beta} = \delta_{\alpha\bar{\beta}} \\ \mathbf{C} &= \begin{pmatrix} 0 & \cdots & 0 & \cdots & 1 \\ \vdots & \ddots & \vdots & \cdot & \vdots \\ 0 & \cdots & 1 & \cdots & 0 \\ \vdots & \cdot & \vdots & \ddots & \vdots \\ 1 & \cdots & 0 & \cdots & 0 \end{pmatrix}. \end{aligned} \tag{13}$$

The annihilation-creation matrix in (5) may be written as

$$\mathbf{K}_{\alpha\beta}^{\delta\gamma} = \mathbf{C}^{\delta\gamma} \mathbf{C}_{\alpha\beta}.$$

2.2 The monodromy matrix

We consider a state with n particles and as is usual in the context of the algebraic Bethe ansatz we define [25, 8] the monodromy matrix by

$$T_{1\dots n,0}(\underline{u}, u_0) = R_{10}(u_{10}) \cdots R_{n0}(u_{n0}) = \begin{array}{c|c|c} & \cdots & \\ \hline & & \\ \hline 1 & n & 0 \end{array} \tag{14}$$

with $\underline{u} = u_1, \dots, u_n$. It is a matrix acting in the tensor product of the “quantum space” $V^{1\dots n} = V_1 \otimes \cdots \otimes V_n$ and the “auxiliary space” V_0 . All vector spaces V_i are isomorphic to a space V whose basis vectors label all kinds of particles. Here $V \cong \mathbb{C}^N$ is the space of the vector representation of $O(N)$.

Suppressing the indices $1 \dots n$ we write the monodromy matrix as (following the notation of Tarasov [15])

$$T_{\alpha}^{\alpha'} = \alpha' \begin{array}{c|c|c} & \cdots & \\ \hline & & \\ \hline \end{array} \alpha = \begin{pmatrix} A_1 & (B_1)_{\hat{\alpha}} & B_2 \\ (C_1)_{\hat{\alpha}'} & (A_2)_{\hat{\alpha}} & (B_3)_{\hat{\alpha}'} \\ C_2 & (C_3)_{\hat{\alpha}} & A_3 \end{pmatrix} \tag{15}$$

where α, α' assume the values $1, 2, \dots, (0), \dots, \bar{2}, \bar{1}$ corresponding to the basis vectors of the auxiliary space $V \cong \mathbb{C}^N$ and $\hat{\alpha}, \hat{\alpha}'$ assume the values $2, \dots, (0), \dots, \bar{2}$ corresponding to the basis vectors of $\hat{V} \cong \mathbb{C}^{N-2}$. We will also use the notation $A = A_1$, $B = B_1$, $C = C_1$ and $D = A_2$ which is an $(N-2) \times (N-2)$ matrix in the auxiliary space. The Yang-Baxter algebra relation for the R-matrix (8) yields

$$T_{1\dots n,a}(\underline{u}, u_a) T_{1\dots n,b}(\underline{u}, u_b) R_{ab}(u_{ab}) = R_{ab}(u_{ab}) T_{1\dots n,b}(\underline{u}, u_b) T_{1\dots n,a}(\underline{u}, u_a) \quad (16)$$

2.3 A lemma

In our approach of the algebraic Bethe ansatz the following lemma replaces commutation rules of the entries of the monodromy matrix. In the conventional approach one derives them from the Yang-Baxter algebra relations (16) and uses them for the algebraic Bethe ansatz.

Lemma 2.2 *For the monodromy matrix the following identity holds*

$$T_{1\dots n,a}(\underline{u}, v) = \mathbf{1}_1 \dots \mathbf{1}_n \mathbf{1}_a + \sum_{i=1}^n c(u_i - v) R_{1a}(u_{1i}) \dots \mathbf{P}_{ia} \dots R_{na}(u_{ni}) + \sum_{j=1}^n d(u_j - v) R_{1a}(\hat{u}_{j1}) \dots \mathbf{K}_{ia} \dots R_{na}(\hat{u}_{jn}) \quad (17)$$

with $\hat{u} = 1/\nu - u$, or in terms of pictures

Proof. The R-matrix $R(u)$ (see (6) and (7)) is meromorphic and has simple poles at $u = 0$ and $u = 1/\nu$ due to the form of $c(u)$ and $d(u)$ such that

$$\begin{aligned} \operatorname{Res}_{v=u_i} T_{1\dots n,a}(\underline{u}, v) &= \operatorname{Res}_{v=u_i} c(u_i - v) R_{1a}(u_{1i}) \dots \mathbf{P}_{ia} \dots R_{na}(u_{ni}) \\ \operatorname{Res}_{v=u_i-1/\nu} T_{1\dots n,a}(\underline{u}, v) &= \operatorname{Res}_{v=u_i-1/\nu} d(u_j - v) R_{1a}(\hat{u}_{j1}) \dots \mathbf{K}_{ia} \dots R_{na}(\hat{u}_{jn}) \end{aligned}$$

holds. The claim follows by Liouville's theorem because $\lim_{v \rightarrow \infty} T_{1\dots n,a}(\underline{u}, v) = \mathbf{1}_1 \dots \mathbf{1}_n \mathbf{1}_a$. ■

Similarly we have for the crossed monodromy matrix

$$T_{a,1\dots n}(v, \underline{u}) = R_{an}(v - u_n) \dots R_{a1}(v - u_1)$$

the relation

$$\begin{aligned} T_{a,1\dots n}(v, \underline{u}) = \mathbf{1}_a \mathbf{1}_1 \dots \mathbf{1}_n + \sum_{i=1}^n c(v - u_i) R_{an}(u_{in}) \dots \mathbf{P}_{ai} \dots R_{a1}(u_{i1}) \\ + \sum_{i=1}^n d(v - u_i) R_{am}(\hat{u}_{mi}) \dots \mathbf{K}_{ai} \dots R_{a1}(\hat{u}_{1i}). \end{aligned} \quad (18)$$

Note that the crossing relation (11) implies

$$T_{a,1\dots n}(v_a, \underline{u}) = \mathbf{C}_{ba} T_{1\dots n,b}(\underline{u}, v_b) \mathbf{C}^{ab} \quad (19)$$

with $v_b = v_a - 1/\nu$.

2.4 The Matrix Π

The nested Bethe ansatz relies on the principle that after each level the rank of the group (or quantum group) is reduced by one. For $SU(N)$ the rank is $N - 1$ and for $O(N)$ it is $[N/2]$. This means that the dimension of the vector representation (where the R-matrix usually acts) is reduced by 1 for the case of $SU(N)$ and by 2 for case of $O(N)$. A more essential difference is that for $SU(N)$ one can use in every level the same R-matrix, because (with a suitable normalization and parameterization) the $SU(N)$ R-matrix does not depend on N . In contrast for $O(N)$ the R-matrix changes after each level, because it depends on N . Therefore we need a new object called matrix Π , which maps the $O(N)$ R-matrix to the $O(N - 2)$ one. We use the notation

$$\begin{aligned} \mathring{R}(u) = R(u, N - 2) = \mathbf{1} + \mathbf{P}c(u) + \mathbf{K}d(u) \\ d(u) = \frac{1}{u - 1/\mathring{\nu}} = \frac{1}{u - 1/\nu + 1} \end{aligned} \quad (20)$$

with $\mathring{\nu} = 2/(N - 4)$. The components of the R-matrix $\mathring{R}(u)$ will be denoted by

$$\mathring{R}_{\alpha\beta}^{\delta\gamma}(u), \quad \alpha, \beta, \gamma, \delta = 2, 3, \dots, (0), \dots, \bar{3}, \bar{2}.$$

In addition to $V^{1\dots m} = V_1 \otimes \dots \otimes V_m$ (1) we introduce

$$\mathring{V}^{1\dots m} = \mathring{V}_1 \otimes \dots \otimes \mathring{V}_m, \quad (21)$$

where the vector spaces $\mathring{V}_i \cong \mathbf{C}^{N-2}$, ($i = 1, \dots, m$) are considered as fundamental (vector) representation spaces of $O(N - 2)$. The space V_i is spanned by the complex basis vectors $|1\rangle, |2\rangle, \dots, |\bar{2}\rangle, |\bar{1}\rangle$ and \mathring{V}_i by $|2\rangle, \dots, |\bar{2}\rangle$.


Definition 2.3 We define the map

$$\Pi_{1\dots m} : V^{1\dots m} \rightarrow \mathring{V}^{1\dots m}$$

recursively by $\Pi_1 = \pi_1$ and

$$\Pi_{1\dots m}(\underline{u}) = (\pi_1 \Pi_{2\dots m}) \bar{e}_a T_{1\dots m, a}(\underline{u}, u_a) \bar{e}^a \quad (22)$$

with the projector $\pi : V \rightarrow \mathring{V} \subset V$, the monodromy matrix (14) and $u_a = u_1 - 1$. The vector $\bar{e}^a \in V_a$ (acting in the auxiliary space of $T_{1\dots m, a}$) and the co-vector $\bar{e}_a \in (V_a)^\dagger$ correspond to the state $\bar{1}$ and have the components $\bar{e}^\alpha = \delta_1^\alpha$, $\bar{e}_\alpha = \delta_\alpha^{\bar{1}}$. This definition may be depicted as



Lemma 2.4 In particular for $m = 2$ the matrix $\Pi_{12}(u_1, u_2)$ may be written as

$$\Pi_{12}(\underline{u}) = \pi_1 \pi_2 + f(u_{12}) \mathring{\mathbf{C}}^{12} \bar{e}_1 e_2 \quad (23)$$

with $e_2 = \mathbf{C}_{2a} \bar{e}^a$ ($e_a = \delta_a^{\bar{1}}$) and $f(u) = -d(1 - u)$. It satisfies the fundamental relation

$$\mathring{R}_{12}(u_{12}) \Pi_{12}(u_1, u_2) = \Pi_{21}(u_2, u_1) R_{12}(u_{12}), \quad (24)$$

where \mathring{R}_{12} is the $O(N - 2)$ R-matrix.

Proof. Equation (23) can be easily derived. We calculate (22) for $m = 2$ with $u_a = u_1 - 1$

$$\begin{aligned} \Pi_{12}(\underline{u}) &= \pi_1 \pi_2 \bar{e}_a T_{12, a}(\underline{u}, u_a) \bar{e}^a \\ &= \pi_1 \pi_2 \bar{e}_a R_{1a}(u_1 - u_a) R_{2a}(u_2 - u_a) \bar{e}^a \\ &= \pi_1 \pi_2 \bar{e}_a (\mathbf{1}_1 \mathbf{1}_a + c(1) \mathbf{P}_{1a}) (\mathbf{1}_2 \mathbf{1}_a + d(u_{21} + 1) \mathbf{K}_{2a}) \bar{e}^a \\ &= \pi_1 \pi_2 + c(1) d(u_{21} + 1) \mathring{\mathbf{C}}^{12} \bar{e}_1 e_2. \end{aligned}$$

Use has been made of $c(1) = -1$ and $\pi_1 \pi_2 \bar{e}_a \mathbf{P}_{1a} \mathbf{K}_{2a} \bar{e}^a = \mathring{\mathbf{C}}^{12} \bar{e}_1 e_2$. Equation (24) is derived for all components. Obviously

$$\mathring{R}_{\hat{\alpha}\hat{\beta}}^{\hat{\beta}'\hat{\alpha}'}(u) \mathring{\mathbf{C}}^{\hat{\alpha}\hat{\beta}} = \mathring{R}_0(u) \mathring{\mathbf{C}}^{\hat{\beta}'\hat{\alpha}'}$$

holds, where the scalar R-matrix eigenvalue is (see (10))

$$\mathring{R}_0(u) = a(u) + (N - 2) \mathring{d}(u).$$

Therefore the relations

$$\left(\overset{\circ}{R}_{12}(u_{12})\Pi_{12}(u_{12})\right)_{\alpha\beta}^{\beta'\alpha'} = \overset{\circ}{R}_{\overset{\circ}{\alpha}\overset{\circ}{\beta}}^{\overset{\circ}{\beta}'\overset{\circ}{\alpha}'}(u_{12})\pi_{\overset{\circ}{\alpha}}^{\overset{\circ}{\alpha}}\pi_{\overset{\circ}{\beta}}^{\overset{\circ}{\beta}} + f(u_{12})\overset{\circ}{R}_0(u_{12})\overset{\circ}{C}^{\beta'\alpha'}\delta_{\overset{\circ}{\alpha}}^{\bar{1}}\delta_{\overset{\circ}{\beta}}^{\bar{1}}$$

and

$$(\Pi_{21}(u_{21})R_{12}(u_{12}))_{\alpha\beta}^{\beta'\alpha'} = \pi_{\overset{\circ}{\beta}'}^{\overset{\circ}{\beta}'}\pi_{\overset{\circ}{\alpha}'}^{\overset{\circ}{\alpha}'}R_{\alpha\beta}^{\beta'\alpha'}(u_{12}) + f(u_{21})\overset{\circ}{C}^{\beta'\alpha'}R_{\alpha\beta}^{\bar{1}\bar{1}}(u_{12})$$

are valid. The claim of the lemma is then equivalent to

$$\begin{aligned} (i) : \overset{\circ}{d}(u) &= d(u) + f(-u)d(u) && \text{for } \alpha \text{ or } \beta \neq 1, \bar{1} \\ (ii) : 0 &= d(u) + f(-u)(1 + d(u)) && \text{for } \alpha = 1, \beta = \bar{1} \\ (iii) : f(u)\overset{\circ}{R}_0(u) &= d(u) + f(-u)(c(u) + d(u)) && \text{for } \alpha = \bar{1}, \beta = 1. \end{aligned}$$

These equations may be easily checked with the amplitudes (7). ■

The matrix Π_{12} may be depicted as

$$\Pi_{\alpha\beta}^{\overset{\circ}{\alpha}\overset{\circ}{\beta}}(u_1, u_2) = \begin{array}{c} \overset{\circ}{\alpha} \\ | \\ \bullet \\ | \\ \alpha \end{array} \begin{array}{c} \overset{\circ}{\beta} \\ | \\ \bullet \\ | \\ \beta \end{array} + f(u_{12}) \begin{array}{c} \overset{\circ}{\alpha} \quad \overset{\circ}{\beta} \\ \diagdown \quad \diagup \\ \bar{1} \quad 1 \\ | \quad | \\ \alpha \quad \beta \end{array}$$

These results can be extended to general m , as presented below.

Lemma 2.5 *The matrix $\Pi_{1\dots m}(\underline{u})$ satisfies*

(a) *in addition to (22) the recursion relation*

$$\Pi_{1\dots m}(\underline{u}) = (\Pi_{1\dots m-1}\pi_m)\bar{e}_b T_{1\dots m,b}(\underline{u}, u_b)\bar{e}^b \quad (25)$$

with $u_b = u_m - 1/\nu + 1$, and

(b) *the fundamental relation*

$$\overset{\circ}{R}_{ij}(u_{ij})\Pi_{\dots ij\dots}(\underline{u}) = \Pi_{\dots ji\dots}(\underline{u})R_{ij}(u_{ij}). \quad (26)$$

(c) *The matrix $\bar{e}_0 T_{1\dots m,0}(\underline{u}, u_0)\bar{e}^0$ acts on $\Pi_{1\dots m}(\underline{u})$ as the unit matrix for arbitrary u_0*

$$\Pi_{1\dots m}(\underline{u})\bar{e}_0 T_{1\dots m,0}(\underline{u}, u_0)\bar{e}^0 = \Pi_{1\dots m}(\underline{u}). \quad (27)$$

(d) *Special components of Π satisfy*

$$\Pi_{1\alpha_2\dots\alpha_m}^{\overset{\circ}{\alpha}_1\dots\overset{\circ}{\alpha}_m}(u_1, \dots, u_m) = 0 \quad (28)$$

$$\Pi_{\overset{\circ}{\alpha}_1\overset{\circ}{\alpha}_2\dots\overset{\circ}{\alpha}_m}^{\overset{\circ}{\alpha}_1\dots\overset{\circ}{\alpha}_m}(u_1, \dots, u_m) = \delta_{\overset{\circ}{\alpha}}^{\overset{\circ}{\alpha}_1} \Pi_{\overset{\circ}{\alpha}_2\dots\overset{\circ}{\alpha}_m}^{\overset{\circ}{\alpha}_2\dots\overset{\circ}{\alpha}_m}(u_2, \dots, u_m) \quad (29)$$

$$\Pi_{\alpha_1\dots\alpha_{m-1}\bar{1}}^{\overset{\circ}{\alpha}_1\dots\overset{\circ}{\alpha}_m}(u_1, \dots, u_m) = 0 \quad (30)$$

$$\Pi_{\alpha_1\dots\alpha_{m-1}\overset{\circ}{\alpha}}^{\overset{\circ}{\alpha}_1\dots\overset{\circ}{\alpha}_m}(u_1, \dots, u_m) = \Pi_{\alpha_1\dots\alpha_{m-1}}^{\overset{\circ}{\alpha}_1\dots\overset{\circ}{\alpha}_{m-1}}(u_1, \dots, u_{m-1})\delta_{\overset{\circ}{\alpha}}^{\overset{\circ}{\alpha}_m}. \quad (31)$$

with $\overset{\circ}{\alpha} \neq 1, \bar{1}$.


The proof of this Lemma is presented in appendix A.

The recursion relations (22) and (25) can be rewritten as (see also lemma 2.4 for $m = 2$)

$$\Pi_{1\dots m}(\underline{u}) = \pi_1 \Pi_{2\dots m} + \sum_{j=2}^m f(u_{1j}) \mathring{R}_{jj-1} \cdots \mathring{R}_{j2} \mathring{C}^{1j} \Pi_{2\dots j\dots m} \bar{e}_1 e_j R_{jm} \cdots R_{jj+1} \quad (32)$$

$$= \Pi_{1\dots m-1} \pi_m + \sum_{j=1}^{m-1} f(u_{jm}) \mathring{R}_{j+1j} \cdots \mathring{R}_{m-1j} \mathring{C}^{jm} \Pi_{1\dots j\dots m-1} \bar{e}_j R_{1j} \cdots R_{j-1j} e_m \quad (33)$$

or in terms of pictures



In particular

$$\Pi_{\bar{1}\alpha_2\dots\alpha_m}^{\hat{\alpha}_1\dots\hat{\alpha}_m}(u_1, \dots, u_m) = \sum_{j=2}^m f(u_{1j}) \left(\mathring{R}_{jj-1} \cdots \mathring{R}_{j2} \mathring{C}^{1j} \Pi_{2\dots j\dots m} e_j R_{jm} \cdots R_{jj+1} \right)_{\alpha_2\dots\alpha_m}^{\hat{\alpha}_1\dots\hat{\alpha}_m} \quad (34)$$

$$\Pi_{\alpha_1\dots\alpha_{m-1}1}^{\hat{\alpha}_1\dots\hat{\alpha}_m}(u_1, \dots, u_m) = \sum_{j=1}^{m-1} f(u_{jm}) \left(\mathring{R}_{j+1j} \cdots \mathring{R}_{m-1j} \mathring{C}^{jm} \Pi_{1\dots j\dots m-1} \bar{e}_j R_{1j} \cdots R_{j-1j} \right)_{\alpha_1\dots\alpha_{m-1}}^{\hat{\alpha}_1\dots\hat{\alpha}_m} \cdot \quad (35)$$

3 The $O(N)$ - difference equation

Let $K_{1\dots n}(\underline{u}) \in V_{1\dots n}$ be a co-vector valued function of $\underline{u} = u_1, \dots, u_n$ with values in $V_{1\dots n}$. The components of this vector are denoted by

$$K_{\alpha_1\dots\alpha_n}(\underline{u}), \quad (\alpha_i = 1, 2, \dots, (0), \dots, \bar{2}, \bar{1}).$$

The following symmetry and periodicity properties of this function are supposed to be valid:

Conditions 3.1

(i) The symmetry property under the exchange of two neighboring spaces V_i and V_j and the variables u_i and u_j , at the same time, is given by

$$K_{\dots ij \dots}(\dots, u_i, u_j, \dots) = K_{\dots ji \dots}(\dots, u_j, u_i, \dots) \tilde{R}_{ij}(u_{ij}), \quad (36)$$

where $\tilde{R}(u) = R(u)/a(u)$ and $R(u)$ is the $O(N)$ R-matrix.

(ii) The **system of matrix difference equations** holds

$$\boxed{K_{1\dots n}(\dots, u'_i, \dots) = K_{1\dots n}(\dots, u_i, \dots) Q_{1\dots n}(\underline{u}; i)}, \quad (i = 1, \dots, n) \quad (37)$$


with $u'_i = u_i + 2/\nu$. The matrix $Q_{1\dots n}(\underline{u}; i) \in \text{End}(V^{1\dots n})$ is defined as the trace

$$Q_{1\dots n}(\underline{u}; i) = \text{tr}_0 \tilde{T}_{Q,1\dots n,0}(\underline{u}, i) \quad (38)$$

of a modified monodromy matrix

$$\tilde{T}_{Q,1\dots n,0}(\underline{u}, i) = \tilde{R}_{10}(u_1 - u'_i) \cdots \mathbf{P}_{i0} \cdots \tilde{R}_{n0}(u_n - u_i).$$

The Yang-Baxter equations for the R-matrix guarantee that these properties are compatible. The shift of $2/\nu$ in eq. (37) could be replaced by an arbitrary κ . For the application to the form factor problem, however, it is fixed to be equal to $2/\nu$ in order to be compatible with crossing symmetry. The properties (i) and (ii) may be depicted as



with the graphical rule that a line changing the "time direction" changes the spectral parameters $u \rightarrow u \pm 1/\nu$ as follows

$$u \downarrow \bigcap_{u-1/\nu} \quad u \uparrow \bigcup_{u+1/\nu}$$

Instead of the Yang-Baxter relation (16) the modified monodromy matrix \tilde{T}_Q satisfies the Zapletal rules [14, 4]. We have for $i = 1, \dots, n$

$$\tilde{T}_Q(\underline{u}; i) T_0(\underline{u}', v) R_{i0}(u_i - v) = R_{i0}(u'_i - v) T_0(\underline{u}, v) \tilde{T}_Q(\underline{u}; i) \quad (39)$$

with $\underline{u}' = u_1, \dots, u'_i, \dots, u_n$ and $u'_i = u_i + 2/\nu$. The $Q_{1\dots n}(\underline{u}; i)$ satisfy the commutation rules

$$Q_{1\dots n}(\dots u_i \dots u_j \dots; i) Q_{1\dots n}(\dots u'_i \dots u_j \dots; j) = Q_{1\dots n}(\dots u_i \dots u_j \dots; j) Q_{1\dots n}(\dots u_i \dots u'_j \dots; i). \quad (40)$$

The following Proposition is obvious

Proposition 3.2 *Let the vector valued function $K_{1\dots n}(\underline{u}) \in V_{1\dots n}$ satisfy (i). Then for all $i = 1, \dots, n$ the relations (3.2) are equivalent to each other and also equivalent to the following periodicity property under cyclic permutation of the spaces and the variables*

$$K_{\alpha_1\alpha_2\dots\alpha_n}(u'_1, u_2, \dots, u_n) = K_{\alpha_2\dots\alpha_n\alpha_1}(u_2, \dots, u_n, u_1). \quad (41)$$

Remark 3.3 *The equations (36,41) imply Watson's equations and crossing relations for the form factors [26].*

Because of proposition 3.2 we mainly consider $Q_{1\dots n}(\underline{u}, i)$ for $i = 1$

$$Q_{1\dots n}(\underline{u}) = \text{tr}_0 \tilde{T}_{Q,1\dots n,0}(\underline{u}) = \prod_{k=2}^n \frac{1}{a(v_{ki})} \text{tr}_0 T_{Q,1\dots n,0}(\underline{u}) \quad (42)$$

with $T_{Q,1\dots n,0}(\underline{u}) = T_{Q,1\dots n,0}(\underline{u}, 1)$. In analogy to eq. (15) we introduce (suppressing the indices $1 \dots n$)

$$T_Q(\underline{u}) \equiv \begin{pmatrix} A_Q(\underline{u}) & B_Q(\underline{u}) & B_{Q,2}(\underline{u}) \\ C_Q(\underline{u}) & D_Q(\underline{u}) & B_{Q,3}(\underline{u}) \\ C_{Q,2}(\underline{u}) & C_{Q,3}(\underline{u}) & A_{Q,3}(\underline{u}) \end{pmatrix}. \quad (43)$$

3.1 The off-shell Bethe ansatz

We will express the co-vector valued function $K_{\underline{\alpha}}(\underline{u})$ in terms of the co-vectors

$$\Psi_{\underline{\alpha}}(\underline{u}, \underline{v}) = L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\alpha}}^{\underline{\beta}}(\underline{u}, \underline{v}) = (L(\underline{v})\Phi(\underline{u}, \underline{v}))_{\underline{\alpha}}, \quad (44)$$

where summation over $\underline{\beta}_1, \dots, \underline{\beta}_m$, $\underline{\beta}_i = 2, \dots, 0, \dots, \bar{2}$ is assumed and $L_{\underline{\beta}}(\underline{v})$ is a co-vector valued function with values in $\dot{V}_{1\dots m} \simeq \mathbb{C}^{N-2} \otimes \dots \otimes \mathbb{C}^{N-2}$. We assume that for $L_{\underline{\beta}}(\underline{v})$ the higher level conditions of 3.1 hold with R and Q replaced by \dot{R} and \dot{Q} (which means N is replaced by $N - 2$)

$$(i)^{(1)} : L_{\dots ij \dots}(\dots, v_i, v_j, \dots) = L_{\dots ji \dots}(\dots, v_j, v_i, \dots) \tilde{R}_{ij}(v_{ij}) \quad (45)$$

$$(ii)^{(1)} : L_{1\dots m}(\dots, v'_i, \dots) = L_{1\dots n}(\dots, v_i, \dots) \dot{Q}_{1\dots m}(\underline{v}, i). \quad (46)$$

with² $v'_i = v_i + 2/\nu$.

The **Bethe ansatz states** are³

$$\Phi_{\underline{\alpha}}^{\underline{\beta}}(\underline{u}, \underline{v}) = \Pi_{\underline{\beta}}^{\underline{\beta}}(\underline{v}) \left(\Omega T_1^{\beta_m}(\underline{u}, v_m) \dots T_1^{\beta_1}(\underline{u}, v_1) \right)_{\underline{\alpha}} = \begin{array}{c} \begin{array}{c} \underline{\beta} \\ \Pi \end{array} \\ \begin{array}{c} \dots \\ \dots \\ \dots \end{array} \\ \begin{array}{c} 1 \\ \dots \\ v_i \\ \dots \\ 1 \end{array} \\ \begin{array}{c} 1 \\ \dots \\ 1 \\ \dots \\ 1 \end{array} \\ \begin{array}{c} u_1 \\ \dots \\ \underline{\alpha} \\ \dots \\ u_n \end{array} \end{array} \cdot \quad (47)$$

²Note that the shift $2/\nu$ is the same in the higher level off-shell Bethe ansatz.

³The $\Phi_{\underline{\alpha}}^{\underline{\beta}}$ are generalizations of the states introduced by Tarasov in [15].

Remark 3.4 The condition (45) implies the symmetry

$$\Psi_{\underline{\alpha}}(\underline{u}, \dots, v_i, v_j, \dots) = \Psi_{\underline{\alpha}}(\underline{u}, \dots, v_j, v_i, \dots). \quad (48)$$

The reference state Ω (“pseudo-vacuum”) is the highest weight co-vector (with weights $w = (n, 0, \dots, 0)$)

$$\Omega_{\underline{\alpha}} = \delta_{\alpha_1}^1 \cdots \delta_{\alpha_n}^1. \quad (49)$$

It satisfies

$$\Omega T(\underline{u}, v) = \Omega \begin{pmatrix} a_1(\underline{u}, v) & 0 & 0 \\ * & a_2(\underline{u}, v) & 0 \\ * & * & a_3(\underline{u}, v) \end{pmatrix}, \quad (50)$$

$$a_1(\underline{u}, v) = \prod_{k=1}^n a(u_k - v), \quad a_2(\underline{u}, v) = 1, \quad a_3(\underline{u}, v) = \prod_{k=1}^n (1 + d(u_k - v)).$$

We also have for $T_Q(\underline{u}) = T_Q(\underline{u}, 1)$

$$\Omega T_Q(\underline{u}) = \Omega \prod_{k=2}^n a(u_{k1}) \begin{pmatrix} 1 & 0 & 0 \\ * & 0 & 0 \\ * & 0 & 0 \end{pmatrix}. \quad (51)$$

The system of difference equations (37) can be solved by means of a nested “off-shell” Bethe ansatz. The first level is given by the **off-shell Bethe ansatz**

$$K_{\underline{\alpha}}(\underline{u}) = \sum_{\underline{v}} g(\underline{u}, \underline{v}) \Psi_{\underline{\alpha}}(\underline{u}, \underline{v}), \quad (52)$$

where the state Ψ is defined by (44) and (47) and the scalar function $g(\underline{u}, \underline{v})$ is

$$g(\underline{u}, \underline{v}) = \prod_{i=1}^n \prod_{j=1}^m \psi(u_i - v_j) \prod_{1 \leq i < j \leq m} \tau(v_i - v_j). \quad (53)$$

The functions $\psi(u)$ and $\tau(v)$ satisfy the functional equations

$$\psi(u') = a(u)\psi(u), \quad \tau(v')a(v') = a(-v)\tau(v). \quad (54)$$

with $u' = u + 2/\nu$ The summation over \underline{v} is specified by

$$\underline{v} = (v_1, \dots, v_m) = (\tilde{v}_1 - 2l_1/\nu, \dots, \tilde{v}_m - 2l_m/\nu), \quad l_i \in \mathbf{Z}, \quad (55)$$

where the \tilde{v}_i are arbitrary constants.

The sums (52) are also called “Jackson-type Integrals” (see e.g. [10] and references therein). Solutions of (54) are

$$\psi(u) = \frac{\Gamma(-\frac{1}{2}\nu + \frac{u}{2}\nu)}{\Gamma(\frac{u}{2}\nu)} \quad (56)$$

$$\tau(v) = v \frac{\Gamma(\frac{1}{2}\nu + \frac{v}{2}\nu)}{\Gamma(1 - \frac{1}{2}\nu + \frac{v}{2}\nu)}. \quad (57)$$

We are now in a position to formulate the main result of this paper.

Theorem 3.5 *Let the co-vector valued function $K_{1\dots n}(\underline{u}) \in V_{1\dots n}$ be given by the Bethe ansatz (52) and let $g(\underline{x}, \underline{u})$ be of the form (53). If in addition the co-vector valued function $L_{1\dots m}(\underline{v}) \in \mathring{V}_{1\dots m}$ satisfies the properties (i)⁽¹⁾ and (ii)⁽¹⁾, i.e. equations (36) and (37) for $O(N-2)$, then $K_{1\dots n}(\underline{u})$ satisfies the equations (36) and (37) for $O(N)$, i.e. $K_{1\dots n}(\underline{u})$ is a solution of the set of difference equations.*

The proof of this theorem can be found in appendix B.

Iterating (52), (44) and theorem 3.5 we obtain the nested off-shell Bethe ansatz with levels $k = 1, \dots, [(N-1)/2] - 1$. The ansatz for level k reads

$$K_{1\dots n_{k-1}}^{(k-1)}(\underline{u}^{(k-1)}) = \sum_{\underline{u}^{(k)}} g^{(k-1)}(\underline{u}^{(k-1)}, \underline{u}^{(k)}) \Psi_{1\dots n_{k-1}}^{(k-1)}(\underline{u}^{(k-1)}, \underline{u}^{(k)}) \quad (58)$$

$$\Psi_{1\dots n_{k-1}}^{(k-1)}(\underline{u}^{(k-1)}, \underline{u}^{(k)}) = (K^{(k)}(\underline{u}^{(k)}) \Phi^{(k-1)}(\underline{u}^{(k-1)}, \underline{u}^{(k)}))_{1\dots n_{k-1}},$$

where $\Phi^{(k)}$ is the Bethe ansatz state (47) and $g^{(k)}$ the function (53) for $O(N-2k)$. The highest levels differ from that of theorem 3.5, but they are given by the $O(3)$ -problem for N odd or the $O(4)$ -problem for N even. These two case are investigated below.

Corollary 3.6 *The system of $O(N)$ matrix difference equations (37) is solved by the nested Bethe ansatz (58) with $K_{1\dots n}(\underline{u}) = K^{(0)}_{1\dots n}(\underline{u})$.*

3.1.1 The off-shell Bethe ansatz for $O(3)$

The $O(3)$ R-matrix is

$$R(u) = \mathbf{1} + c(u)P + d(u)K, \quad c(u) = \frac{-1}{u}, \quad d(u) = \frac{1}{u-1/2}.$$

The solution of the difference equations (36)-(38) is again given by the off-shell Bethe ansatz (52)-(57). The Bethe vector Ψ is expressed in terms of the co-vectors (47)

$$\Psi_{\underline{\alpha}}(\underline{u}, \underline{v}) = L(\underline{v}) \Phi_{\underline{\alpha}}(\underline{u}, \underline{v}),$$

where the scalar function $L(\underline{v})$ has to satisfy

$$\begin{aligned} L(\dots, v_i, v_j, \dots) &= L(\dots, v_j, v_i, \dots) \tilde{R}(v_{ij}) \\ L(v'_1, v_2, \dots, v_m) &= L(v_2, \dots, v_m, v_1) \end{aligned} \quad (59)$$

with $\tilde{R}(v) = \frac{(v+1)(v-1/2)}{(v-1)(v+1/2)}$. For $N = 3$ i.e. $\nu = 2$ we have

$$\psi(u) = \frac{1}{u-1}, \quad \tau(v) = v^2.$$

and $v' = v + 1$. The the minimal solution of the equations (59) is

$$\begin{aligned} L(\underline{v}) &= \prod_{1 \leq i < j \leq m} L(v_{ij}) \\ L(v) &= \frac{\pi(v-1/2)}{4v(v-1)} \tan \pi v. \end{aligned}$$

The $O(3)$ weight of the state $\Psi_{\underline{\alpha}}(\underline{u}, \underline{v})$ is

$$w = n - m. \quad (60)$$

3.1.2 The off-shell Bethe ansatz for $O(4)$

The $O(4)$ R-matrix is

$$R(u) = \mathbf{1} + c(u)P + d(u)K, \quad c(u) = \frac{-1}{u}, \quad d(u) = \frac{1}{u-1}.$$

We could apply theorem 3.5 and write the off-shell Bethe ansatz for $O(4)$ in terms of an $O(2)$ problem. However, the latter cannot be solved by the Bethe ansatz because the R-matrix is diagonal (note that $R_{1\bar{1}}^{\bar{1}1} = 0$). But there is another way to solve the $O(4)$ problem. The group isomorphism $O(4) \simeq SU(2) \otimes SU(2)$ reflects itself in terms of the corresponding R-matrices. Indeed, the $O(4)$ R-matrix can be written as a tensor product of two $SU(2)$ R-matrices, or more precisely

$$\left(\tilde{R}^{O(4)}\right)_{\alpha\beta}^{\delta\gamma} \Gamma_{AB}^{\alpha} \Gamma_{CD}^{\beta} = \Gamma_{C'D'}^{\delta} \Gamma_{A'B'}^{\gamma} \left(\tilde{R}_{+}^{SU(2)}\right)_{AC}^{C'A'} \left(\tilde{R}_{-}^{SU(2)}\right)_{BD}^{D'B'}$$

The $SU(2)$ R-matrices $\tilde{R}_{\pm}^{SU(2)}(u) = R_{\pm}^{SU(2)}(u)/a(u)$ correspond to the spinor representations of $O(4)$ with positive (negative) chirality

$$R_{\pm}^{SU(2)} = \mathbf{1} + c(u)\mathbf{P},$$

where the amplitude $c(u) = -1/u$ is again given by (7). The relative R-matrix for states of different chirality is trivial $\tilde{R} = \mathbf{1}$. The intertwiners are

$$\Gamma_{AB}^{\alpha} = (\gamma_{+}\gamma^{\alpha}\mathbf{C})_{AB}$$

with the $O(4)$ gamma matrices γ^{α} , $\gamma_{+} = \frac{1}{2}(1 + \gamma^5)$ and the charge conjugation matrix \mathbf{C} . For more details see [27, 28]. In the complex basis of the $O(4)$ and the fundamental representations of the $SU(2)$ the states have the $O(4)$ weights

vector states	$O(4)$ weights	spinor states	$O(4)$ weights
1	(1, 0)	\uparrow_{+}	$(\frac{1}{2}, \frac{1}{2})$
2	(0, 1)	\downarrow_{+}	$(-\frac{1}{2}, -\frac{1}{2})$
$\bar{2}$	(0, -1)	\uparrow_{-}	$(\frac{1}{2}, -\frac{1}{2})$
$\bar{1}$	(-1, 0)	\downarrow_{-}	$(-\frac{1}{2}, \frac{1}{2})$.

(61)

Because of weight conservation the intertwiner matrix is diagonal in this basis and is calculated to be

$$\Gamma_{AB}^\alpha = \begin{pmatrix} \Gamma_{\uparrow\uparrow\uparrow-}^1 & 0 & 0 & 0 \\ 0 & \Gamma_{\uparrow\uparrow\downarrow-}^2 & 0 & 0 \\ 0 & 0 & \Gamma_{\downarrow\uparrow\uparrow-}^{\bar{2}} & 0 \\ 0 & 0 & 0 & \Gamma_{\downarrow\uparrow\downarrow-}^{\bar{1}} \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (62)$$

We also use the dual intertwiner Γ_α^{AB} with

$$\sum_{A,B} \Gamma_{AB}^{\alpha'} \Gamma_\alpha^{AB} = \delta_\alpha^{\alpha'}, \quad \sum_\alpha \Gamma_\alpha^{A'B'} \Gamma_{AB}^\alpha = \delta_A^{A'} \delta_B^{B'}. \quad (63)$$

We write the co-vector valued function $K_\alpha(\underline{u})$ as

$$K_\alpha(\underline{u}) = K_{\underline{A}}^{(+)}(\underline{u}) K_{\underline{B}}^{(-)}(\underline{u}) \Gamma_\alpha^{AB} \quad (64)$$

$$\begin{array}{c} \text{---} K \text{---} \\ | \\ \underline{\alpha} \end{array} = \begin{array}{c} \text{---} K^{(+)} \text{---} \quad \text{---} K^{(-)} \text{---} \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \underline{A} \quad \underline{B} \\ | \\ \underline{\alpha} \end{array}, \quad (65)$$

where $\Gamma_\alpha^{AB} = \prod_{i=1}^n \Gamma_{\alpha_i}^{A_i B_i}$. The transfer matrix $\text{tr} \tilde{T}^{O(4)}(\underline{u}, v)$ can also be decomposed such that

$$\begin{aligned} K_\alpha(\underline{u}) \left(\tilde{T}^{O(4)} \right)_\gamma^\gamma(\underline{u}, v) &= K_\alpha(\underline{u}) \Gamma_{A'B'}^\gamma \left(\tilde{T}_+^{SU(2)} \right)_A^{A'}(\underline{u}, v) \left(\tilde{T}_-^{SU(2)} \right)_B^{B'}(\underline{u}, v) \Gamma_\gamma^{AB} \\ &= \left(K_{\underline{A}}^{(+)}(\underline{u}) \left(\tilde{T}_+^{SU(2)} \right)_A^A(\underline{u}, v) \right) \left(K_{\underline{B}}^{(-)}(\underline{u}) \left(\tilde{T}_-^{SU(2)} \right)_B^B(\underline{u}, v) \right) \Gamma_\alpha^{AB} \end{aligned}$$

$$\begin{array}{c} \text{---} K \text{---} \\ | \quad | \\ \gamma \quad \gamma \end{array} = \begin{array}{c} \text{---} K^{(+)} \text{---} \quad \text{---} K^{(-)} \text{---} \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ A \quad A \quad B \quad B \\ | \quad | \end{array},$$

where (63) has been used. Therefore $K_\alpha(\underline{u})$ satisfies the $O(4)$ symmetry relation (36) and the difference equation (37) if the $K_{\underline{A}}^{(\pm)}(\underline{u})$ satisfy the corresponding $SU(2)$ relations.

The $SU(2)$ on-shell Bethe ansatz is well known and the off-shell case has been solved in [14, 6, 29]

$$K_{\underline{A}}(\underline{u}) = \sum_{\underline{v}} g(\underline{u}, \underline{v}) \Psi_{\underline{A}}(\underline{u}, \underline{v}) \quad (66)$$

$$\Psi(\underline{u}, \underline{v}) = \Omega C(\underline{u}, v_m) \dots C(\underline{u}, v_1), \quad (67)$$

where $\sum_{\underline{v}}$ and $g(\underline{u}, \underline{v})$ are given by (53)-(57). For $N = 4$ i.e. $\nu = 1$ we have

$$\psi(u) = \frac{\Gamma(-\frac{1}{2} + \frac{u}{2})}{\Gamma(\frac{u}{2})}, \quad \tau(v) = v.$$

The $SU(2)$ weights of the state (67) are $w = (n - m, m)$ and due to (61) the $O(4)$ weights are

$$w = \begin{cases} (n - m, -m) & \text{for positive chirality spinors} \\ (n - m, m) & \text{for negative chirality spinors.} \end{cases}$$

Therefore the $O(4)$ weights of (64) are (see also [17])

$$w = (n - n_- - n_+, n_- - n_+), \quad (68)$$

where n_{\pm} are the numbers of positive (negative) chirality C -operators.

4 Weights of off-shell $O(N)$ Bethe vectors

In this section we analyze some group theoretical properties of off-shell Bethe states. We show that they are highest weight states and we calculate the weights. The first result is not only true for the conventional Bethe ansatz, which solves an eigenvalue problem and which is well known, but it is also true, as we will show, for the off-shell one which solves a difference equation (or a differential equation).

By the asymptotic expansion of the R-matrix (5) and the monodromy matrix (14) we get for $u \rightarrow \infty$

$$R_{ab}(u) = \mathbf{1}_{ab} - \frac{1}{u} M_{ab} + O(u^{-2}) \quad (69)$$

$$M_{ab} = \mathbf{P}_{ab} - \mathbf{K}_{ab}. \quad (70)$$

More explicitly eq. (14) gives

$$T_{1\dots n,a}(\underline{u}, u) = \mathbf{1}_{1\dots n,a} + \frac{1}{u} M_{1\dots n,a} + O(u^{-2}) \quad (71)$$

$$M_{1\dots n,a} = (\mathbf{P}_{1a} - \mathbf{K}_{1a}) + \dots + (\mathbf{P}_{na} - \mathbf{K}_{na}). \quad (72)$$

The matrix elements of $M_{1\dots n,a}$, as a matrix in the auxiliary space, are the $O(N)$ Lie algebra generators. In the following we will consider only operators acting in the fixed tensor product space $V^{1\dots n}$ of (1). Therefore we will omit the indices $1\dots n$. In terms of the matrix elements in the auxiliary space V_a the generators act on the basis states as

$$\langle \alpha_1, \dots, \alpha_i, \dots, \alpha_n | M_{\alpha}^{\alpha'} = \sum_{i=1}^n \left(\delta_{\alpha\alpha_i} \langle \alpha_1, \dots, \alpha', \dots, \alpha_n | - \delta_{\alpha'\bar{\alpha}_i} \langle \alpha_1, \dots, \bar{\alpha}, \dots, \alpha_n | \right). \quad (73)$$

The diagonal elements of $M_{\alpha}^{\alpha'}$ are the weight operators W_{α} with

$$\langle \alpha_1, \dots, \alpha_i, \dots, \alpha_n | W_{\alpha} = w_{\alpha} \langle \alpha_1, \dots, \alpha_i, \dots, \alpha_n |$$

$$w_{\alpha} = n_{\alpha} - n_{\bar{\alpha}}$$

where n_{α} is the number of particles α in the state. It is sufficient to consider only the weights

$$w = (w_1, \dots, w_{[N/2]}) \quad (74)$$

because of $W_\alpha = -W_{\bar{\alpha}}$ and $\langle \underline{\alpha} | W_0 = 0$ for N odd.

The Yang-Baxter relations (16) yield for $u_a \rightarrow \infty$

$$[M_a + M_{ab}, T_b(u_b)] = 0 \quad (75)$$

and if additionally $u_b \rightarrow \infty$, we get

$$[M_a + M_{ab}, M_b] = 0, \quad (76)$$

or for the matrix elements (in the real basis)

$$[M_\alpha^{\alpha'}, T_\beta^{\beta'}(u)] = -\delta_\alpha^{\beta'} T_\beta^{\alpha'}(u) + \delta^{\alpha'\beta'} T_\beta^\alpha(u) + T_\alpha^{\beta'}(u) \delta_\beta^{\alpha'} - T_{\alpha'}^{\beta'}(u) \delta_{\alpha\beta} \quad (77)$$

$$[M_\alpha^{\alpha'}, M_\beta^{\beta'}] = -\delta_\alpha^{\beta'} M_\beta^{\alpha'} + \delta^{\alpha'\beta'} M_\beta^\alpha + M_\alpha^{\beta'} \delta_\beta^{\alpha'} - M_{\alpha'}^{\beta'} \delta_{\alpha\beta}. \quad (78)$$

Equation (78) represents the structure relations of the $O(N)$ Lie algebra and (77) the $O(N)$ -covariance of T . In particular the transfer matrix is invariant

$$[M_\alpha^{\alpha'}, \text{tr } T(u)] = 0. \quad (79)$$

Theorem 4.1 1. *If the co-vector valued function*

$$K_{\underline{\alpha}}(\underline{u}) = \sum_{\underline{v}} g(\underline{u}, \underline{v}) L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\alpha}}^{\underline{\beta}}(\underline{u}, \underline{v})$$

is given by the nested off-shell Bethe ansatz (58) the weights (74) are $w =$

$$(w_1, \dots, w_{[N/2]}) = \begin{cases} (n - n_1, \dots, n_{[N/2]-1} - n_{[N/2]}) & \text{for } N \text{ odd} \\ (n - n_1, \dots, n_{[N/2]-2} - n_-, n_- - n_+, n_- - n_+) & \text{for } N \text{ even.} \end{cases}$$

2. *If $K_{\underline{\alpha}}(\underline{u})$ satisfies the conditions of theorem 3.5 and if $L_{\underline{\beta}}(\underline{v})$ is a highest weight state, then $K_{\underline{\alpha}}(\underline{u})$ is a highest weight state:*

$$K(\underline{u}) M_\alpha^{\alpha'} = 0 \text{ for } \alpha' < \alpha.$$

(Recall that $\alpha' < \alpha$ is to be understood corresponding to the ordering $1, 2, \dots, [N/2], (0), \overline{[N/2]}, \dots, \bar{2}, \bar{1}$.)

3. *The weights satisfy the highest weight condition*

$$\begin{cases} w_1 \geq w_2 \geq \dots \geq w_{[N/2]} \geq 0 & \text{for } N \text{ odd} \\ w_1 \geq w_2 \geq \dots \geq |w_{[N/2]}| & \text{for } N \text{ even.} \end{cases}$$

The proof of this theorem can be found in appendix C. We mention that for N even the highest weight property was already discussed in appendix B of [17].

5 Conclusion

In this article we solved the $O(N)$ -matrix difference equations by means of the off-shell algebraic nested Bethe ansatz. We introduced a new object called Π -matrix to overcome the difficulties connected to the special peculiarities of the $O(N)$ symmetric R-matrix structure. The highest weights properties of the solutions were analyzed. We believe that our construction can also be applied to the cases with similar group theoretical complexities, such as B_n, C_n, D_n Lie algebras and superalgebra $Osp(n|2m)$ (see [16]).

Acknowledgment: The authors have profited from discussions with A. Fring, R. Schrader, F. Smirnov and A. Belavin. H.B. thanks R. Flume, R. Poghossian and P. Wiegmann for valuable discussions. H.B. and M.K. were supported by Humboldt Foundation and H.B. is also supported by the Armenian grant 11-1_c028. A.F. acknowledges support from DAAD (Deutscher Akademischer Austausch Dienst) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico).

Appendix

A Proof of Lemma 2.5

Proof.


(a) We prove (25) by induction: It is true for $m = 2$, because similar to (23)

$$\pi_1 \pi_2 \bar{e}_b T_{12,b}(\underline{u}, u_b) \bar{e}^b = \pi_1 \pi_2 + f(u_{12}) \mathring{C}^{12} \bar{e}_1 e_2.$$

We assume (25) for $m - 1$ and replace in the definition (22) $\Pi_{2\dots m}$ as given by (25)

$$\begin{aligned} \Pi_{1\dots m} &= \pi_1 \Pi_{2\dots m} \bar{e}_a T_{1\dots m,a} \bar{e}^a \\ &= \pi_1 \left(\Pi_{2\dots m-1} \pi_m \bar{e}_b T_{2\dots m,b} \bar{e}^b \right) \bar{e}_a T_{1\dots m,a} \bar{e}^a \\ &= (\pi_1 \Pi_{2\dots m-1} \pi_m) \bar{e}_b T_{1\dots m,b} \bar{e}^b \bar{e}_a T_{1\dots m,a} \bar{e}^a \\ &= (\pi_1 \Pi_{2\dots m-1} \pi_m) \bar{e}_a T_{1\dots m,a} \bar{e}^a \bar{e}_b T_{1\dots m,b} \bar{e}^b \\ &= (\pi_1 \Pi_{2\dots m-1} \pi_m) \bar{e}_a T_{1\dots m-1,a} \bar{e}^a \bar{e}_b T_{1\dots m,b} \bar{e}^b \\ &= (\Pi_{1\dots m-1} \pi_m) \bar{e}_b T_{1\dots m,b} \bar{e}^b. \end{aligned}$$

for $u_a = u_1 - 1$ and $u_b = u_m - 1/\nu + 1$. Going from equality 2 to equality 3



we have $T_{2\dots m,b}$ replaced by $T_{1\dots m,b}$, which means $R_{1b}(u_{1b})$ may be replaced by $\mathbf{1}_{1b}$, because

$$\bar{e}_b R_{1b}(u_{1b}) \bar{e}_a R_{1a}(u_{1a}) = \bar{e}_b \bar{e}_a R_{1a}(1) + c(u_{1b}) \bar{e}_1 \bar{e}_a R_{1a}(1) = \bar{e}_b \bar{e}_a R_{1a}(1)$$

holds, where $\bar{e}_1 \bar{e}_a R_{1a}(1) = \bar{e}_1 \bar{e}_a a(1) = 0$ has been used. Similarly, equality 5 holds. Equality 4 holds because the Yang-Baxter equation for R implies that $\bar{e}_a T_{1\dots m,a} \bar{e}^a$ and $\bar{e}_b T_{1\dots m,b} \bar{e}^b$ commute.

- (b) Again we prove (26) by induction. For $m = 2$ the claim was proved in section 2.4, for $m > 2$ it follows for $1 < i < j$ from (22) and for $i < j < m$ from (25).
- (c) The proof of equation (27) is similar to that of a). We commute $T(u_a)$ and $T(u_0)$, use $\bar{e}_0 \pi_1 R_{10}(u_{10}) R_{1a}(1) = \bar{e}_0 \pi_1$ and apply induction:

$$\begin{aligned} \Pi_{1\dots m} \bar{e}_0 T_{1\dots m,0}(u_0) \bar{e}^0 &= (\pi_1 \Pi_{2\dots m}) \bar{e}_a T_{1\dots m,a} \bar{e}^a \bar{e}_0 T_{1\dots m,0}(u_0) \bar{e}^0 \\ &= (\pi_1 \Pi_{2\dots m}) \bar{e}_0 T_{1\dots m,0}(u_0) \bar{e}^0 \bar{e}_a T_{1\dots m,a} \bar{e}^a \\ &= \pi_1 (\Pi_{2\dots m}) \bar{e}_0 T_{2\dots m,0}(u_0) \bar{e}^0 \bar{e}_a T_{1\dots m,a} \bar{e}^a \\ &= (\pi_1 \Pi_{2\dots m}) \bar{e}_a T_{1\dots m,a} \bar{e}^a = \Pi_{1\dots m} . \end{aligned}$$

- (d) Equations (28) and (29) follow from (22) and $R_{1\alpha}^{\bar{1}\hat{\alpha}1}(1) = 0$, $R_{\hat{\alpha}\alpha}^{\bar{1}\hat{\alpha}1}(1) = \delta_{\hat{\alpha}}^{\hat{\alpha}1} \delta_{\alpha}^{\bar{1}}$ and analogously (30) and (31).

■

B Proof of the main theorem 3.5

In the following we use the convention that α, β etc. take the values $1, 2, \dots, (0), \dots, \bar{2}, \bar{1}$ and $\hat{\alpha}, \hat{\beta}$ etc. take the values $2, \dots, (0), \dots, \bar{2}$.

$$\begin{aligned} K_{\underline{\alpha}}(\underline{u}) &= \sum_{\underline{v}} g(\underline{u}, \underline{v}) \Psi_{\underline{\alpha}}(\underline{u}, \underline{v}), \quad \Psi_{\underline{\alpha}}(\underline{u}, \underline{v}) = L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\alpha}}^{\hat{\beta}}(\underline{u}, \underline{v}) \\ \Phi_{\underline{\alpha}}^{\hat{\beta}}(\underline{u}, \underline{v}) &= \Pi_{\underline{\beta}}^{\hat{\beta}}(\underline{v}) \left(\Omega T_1^{\beta m}(\underline{u}, v_m) \dots T_1^{\beta 1}(\underline{u}, v_1) \right)_{\underline{\alpha}} . \end{aligned}$$

(i) **Proof.** Property (i) in the form of (36) follows directly from the Yang-Baxter equations and the action of the R-matrix on the pseudo-ground state Ω


$$\begin{aligned} (T_{\dots j i \dots})_1^{\hat{\beta}}(\dots u_j, u_i \dots) R_{ij}(u_{ij}) &= R_{ij}(u_{ij}) (T_{\dots i j \dots})_1^{\hat{\beta}}(\dots u_i, u_j \dots) \\ \Omega_{\dots i j \dots} R_{ij}(u_{ij}) &= a(u_{ij}) \Omega_{\dots i j \dots} . \end{aligned}$$

■

(ii) **Proof.** We prove

$$K_{1\dots n}(\underline{u}') = K_{1\dots n}(\underline{u}) Q_{1\dots n}(\underline{u}), \quad (\text{B.1})$$

where $\underline{u}' = (u_1 + 2/\nu, u_2, \dots, u_n)$. The matrix $Q(\underline{u}) = Q(\underline{u}, 1)$ is given by (42). Note that we assign to the auxiliary space of $\tilde{T}_Q(\underline{u})$ the spectral parameter u_1 on the right hand side and $u'_1 = u_1 + 2/\nu$ on the left hand side. The difference equation (B.1) may be depicted as



where we use the rule that the rapidity of a line changes by $2/\nu$ if the line bends by 360° in the positive sense. In the following we will suppress the indices $1 \dots n$. We are now going to prove (B.1) in the form

$$K(\underline{u}) \left(A_Q(\underline{u}) + D_{Q\beta}^{\dot{\beta}}(\underline{u}) + A_{3,Q}(\underline{u}) \right) = \prod_{k=2}^n a(u_{k1}) K(\underline{u}'), \quad (\text{B.2})$$

where $K(\underline{u})$ is a co-vector valued function as given by eq. (52) and the Bethe ansatz state (44). To analyze the left hand side of eq. (B.2) we proceed as follows: We apply the trace of T_Q to the co-vector $K(\underline{u})$. In particular we calculate $\Phi_{\underline{\beta}}^{\dot{\beta}}(\underline{u}, \underline{v}) T_Q(\underline{u})$

$$\Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(\underline{u}, v_m) \cdots T_1^{\beta_1}(\underline{u}, v_1) (T_Q(\underline{u}))_{\gamma}^{\gamma'} =$$

We now proceed as usual in the algebraic Bethe ansatz and push $A_Q(\underline{u})$, $D_{Q\beta}^{\dot{\beta}}(\underline{u})$ and $A_{3,Q}(\underline{u})$ through all the $T_1^{\beta_i}$ -operators. As usual we obtain wanted terms and unwanted terms. We first find that the wanted contribution from $A_Q(\underline{u})$ already gives the result we are looking for. Secondly the wanted contributions from $D_{Q\beta}^{\dot{\beta}}(\underline{u})$ and $A_{3,Q}(\underline{u})$ applied to Ω give zero because of (51). Thirdly the unwanted contributions from $A_Q(\underline{u})$, $D_{Q\beta}^{\dot{\beta}}(\underline{u})$ and $A_{3,Q}(\underline{u})$ can be written as differences which cancel after summation over the v_j . All these three facts can be seen as follows.

The “wanted terms” from A_Q are obtained if one writes the Zapletal commutation rule (39) as

$$T_1^{\beta_k}(\underline{u}, v_k) A_Q(\underline{u}) = A_Q(\underline{u}) T_1^{\beta_i}(\underline{u}', v_k) a(u_1 - v_k) + uw.$$

Therefore we obtain

$$(L(\underline{v}) \Pi_{\underline{\beta}} \underline{u}, \underline{v})_{\underline{\beta}} \Omega T_1^{\beta_m}(\underline{u}, v_m) \cdots T_1^{\beta_1}(\underline{u}, v_1) A_Q(\underline{u}) = w^A + uw^A$$

with

$$\begin{aligned} w^A(\underline{u}, \underline{v}) &= (L(\underline{v})\Pi\underline{u}, \underline{v})_{\underline{\beta}} \Omega A_Q(\underline{u}) T_1^{\beta_m}(\underline{u}', v_m) \cdots T_1^{\beta_1}(\underline{u}', v_1) \prod_{k=1}^m a(u_1 - v_k) \\ &= \prod_{k=2}^n a(u_{k1}) \prod_{k=1}^m a(u_1 - v_k) L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\beta}}^{\dot{\beta}}(\underline{u}', \underline{v}) \end{aligned} \quad (\text{B.3})$$

and similarly $w^D = w^{A_3} = 0$, because of (51). Inserted into (52) this yields

$$\begin{aligned} (K(\underline{u})Q)^{\text{wanted}}(\underline{u}) &= \sum_{\underline{v}} g(\underline{u}, \underline{v}) \prod_{k=1}^m a(u_1 - v_k) L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\beta}}^{\dot{\beta}}(\underline{u}', \underline{v}) \\ &= \sum_{\underline{v}} g(\underline{u}', \underline{v}) L_{\underline{\beta}}(\underline{v}) \Phi_{\underline{\beta}}^{\dot{\beta}}(\underline{u}', \underline{v}) = K(\underline{u}') \end{aligned}$$

because

$$g(\underline{u}, \underline{v}) \prod_{k=1}^m a(u_1 - v_k) = g(\underline{u}', \underline{v}),$$

where (54) has been used. Therefore it remains to prove that the unwanted terms cancel. This will follow from the lemma below. ■

We apply $Q_{1\dots n}(\underline{u})$ to the state $\Psi_{1\dots n}(\underline{u}, \underline{v})$ (suppressing the quantum space indices)

$$\Psi(\underline{u}, \underline{v}) \left(A_Q(\underline{u}) + D_{Q_{\underline{\beta}}^{\dot{\beta}}}(\underline{u}) + A_{3,Q}(\underline{u}) \right) = \text{wanted} + \text{unwanted}.$$

The wanted contribution has been calculated above and the unwanted terms may be written as (see subsection B.1)

$$\begin{aligned} \text{unwanted} &= uw^A + uw^D + uw^{A_3} \\ uw^A &= \sum_{i=1}^m \left(uw_C^{A,i} \right)_{\dot{\gamma}} C_Q^{\dot{\gamma}} + \sum_{i=1}^m \sum_{\substack{j=1 \\ j \neq i}}^m uw_{C_2}^{A,ij} C_{2,Q} \\ uw^D &= \sum_{i=1}^m \left(uw_C^{D,i} \right)_{\dot{\gamma}} C_Q^{\dot{\gamma}} + \sum_{i=1}^m \sum_{\substack{j=1 \\ j \neq i}}^m uw_{C_2}^{D,ij} C_{2,Q} + \sum_{i=1}^m \left(uw_{C_3}^{D,i} \right)_{\dot{\gamma}} (C_{3,Q})_{\dot{\gamma}'} \dot{C}^{\dot{\gamma}'\dot{\gamma}} \\ uw^{A_3} &= \sum_{i=1}^m \sum_{\substack{j=1 \\ j \neq i}}^m uw_{C_2}^{A_3,ij} C_{2,Q} + \sum_{i=1}^m \left(uw_{C_3}^{A_3,i} \right)_{\dot{\gamma}} (C_{3,Q})_{\dot{\gamma}'} \dot{C}^{\dot{\gamma}'\dot{\gamma}}. \end{aligned}$$

Lemma B.1 *The unwanted terms satisfy the relations*

$$\left(uw_C^{D,i} \right)_{\dot{\gamma}}(\underline{u}, \underline{v}) g(\underline{u}, \underline{v}) = - \left(uw_C^{A,i} \right)_{\dot{\gamma}}(\underline{u}, \underline{v}^{(i)}) g(\underline{u}, \underline{v}^{(i)}) \quad (\text{B.4})$$

$$\left(uw_{C_3}^{A_3,i} \right)_{\dot{\gamma}}(\underline{u}, \underline{v}) g(\underline{u}, \underline{v}) = - \left(uw_{C_3}^{D,i} \right)_{\dot{\gamma}}(\underline{u}, \underline{v}^{(i)}) g(\underline{u}, \underline{v}^{(i)}) \quad (\text{B.5})$$

$$uw_{C_2}^{D,ij}(\underline{u}, \underline{v}^{(j)}) g(\underline{u}, \underline{v}^{(j)}) = -uw_{C_2}^{A,ij}(\underline{u}, \underline{v}^{(ij)}) g(\underline{u}, \underline{v}^{(ij)}) - uw_{C_2}^{A_3,ij}(\underline{u}, \underline{v}) g(\underline{u}, \underline{v}) \quad (\text{B.6})$$

with the notation

$$\begin{aligned}\underline{v} &= v_1, \dots, v_m \\ \underline{v}^{(i)} &= v_1, \dots, v'_i, \dots, v_m \\ \underline{v}^{(ij)} &= v_1, \dots, v'_i, \dots, v'_j, \dots, v_m \\ v' &= v + 2/\nu.\end{aligned}$$

Equations (B.4) - (B.6) imply that all unwanted terms cancel after summation over the \underline{v} in the Jackson-type Integral (52).

Proof. The first relation (B.4) is the same as the relation of the unwanted terms for $SU(N)$. The two others are new and more complicated to derive. We can calculate the following unwanted contributions explicitly (see subsection B.1)

$$\left(uw_{C_1}^{A,i}\right)_{\tilde{\gamma}}(\underline{u}, \underline{v}) = -c(u'_1 - v_i)X_{\tilde{\gamma}}^{(i)}(\underline{u}, \underline{v}) \quad (\text{B.7})$$

$$\left(uw_{C_1}^{D,i}\right)_{\tilde{\gamma}}(\underline{u}, \underline{v}) = c(u_1 - v_i)X_{\tilde{\gamma}}^{(i)}(\underline{u}, \underline{v}^{(i)})\chi_i(\underline{u}, \underline{v}) \quad (\text{B.8})$$

$$\left(uw_{C_3}^{D,i}\right)_{\tilde{\gamma}}(\underline{u}, \underline{v}) = -f(u'_1 - v_i)X_{\tilde{\gamma}}^{(i)}(\underline{u}, \underline{v}) \quad (\text{B.9})$$

$$\left(uw_{C_3}^{A_3,i}\right)_{\tilde{\gamma}}(\underline{u}, \underline{v}) = f(u_1 - v_i)X_{\tilde{\gamma}}^{(i)}(\underline{u}, \underline{v}^{(i)})\chi_i(\underline{u}, \underline{v}) \quad (\text{B.10})$$

with

$$X_{\tilde{\gamma}}^{(i)}(\underline{u}, \underline{v}) = L(v_i, \underline{v}_i)_{\tilde{\gamma}\tilde{\beta}_i} \Phi^{\tilde{\beta}_i}(\underline{u}, \underline{v}_i) \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik})a_1(\underline{u}, v_i) \quad (\text{B.11})$$

and

$$\chi_i(\underline{u}, \underline{v}) = \frac{a_2(\underline{u}, v_i)}{a_1(\underline{u}, v'_i)} \prod_{\substack{k=1 \\ k \neq i}}^m \frac{a(v_{ki})}{a(v'_{ik})}, \quad (\text{B.12})$$

where a_1 and a_2 are defined in (50). We use the short notations \underline{v}_i and $\tilde{\beta}_i$ which means that v_i and $\tilde{\beta}_i$ are missing, respectively. The remaining unwanted terms are

$$uw_{C_2}^{A,ij}(\underline{u}, \underline{v}) = -c(u'_1 - v_i)X^{(ij)}(\underline{v}) \quad (\text{B.13})$$

$$uw_{C_2}^{D,ij}(\underline{u}, \underline{v}) = (c(u_1 - v_i) + f(u'_1 - v_j))X^{(ij)}(\underline{v}^{(i)})\chi_i(\underline{u}, \underline{v}) \quad (\text{B.14})$$

$$uw_{C_2}^{A_3,ij}(\underline{u}, \underline{v}) = -f(u_1 - v_j)X^{(ij)}(v'_i, v'_j, \underline{v}_{ij})\chi_i(\underline{u}, \underline{v}^{(j)})\chi_j(\underline{u}, \underline{v}) \quad (\text{B.15})$$

with

$$X^{(ij)}(\underline{u}, \underline{v}) = f(v_{ij})a(v_{ij})L(v_i, v_j, \underline{v}_{ij})\tilde{\mathbf{C}}^{ij}\Phi(\underline{u}, \underline{v}_{ij}) \prod_{\substack{k=1 \\ k \neq i,j}}^m (a(v_{ik})a(v_{jk}))a_1(v_i)a_1(v_j), \quad (\text{B.16})$$

where again \underline{v}_{ij} means that v_i and v_j are missing. The claims (B.4) - (B.6) follow from the shift property of the function $g(\underline{u}, \underline{v})$ defined in (53)

$$\chi_i(\underline{u}, \underline{v})g(\underline{u}, \underline{v}) = g(\underline{u}, \underline{v}^{(i)})$$

which is due to the shift properties (54) of the functions $\psi(v)$ and $\tau(v)$.

The states $X_{\dot{\gamma}}^{(1)}$ and $X^{(12)}$ may be depicted as

$$X_{\dot{\gamma}}^{(1)}(\underline{u}, \underline{v}) = \begin{array}{c} \text{---} L \text{---} \\ | \\ \text{---} \Pi \text{---} \\ | \\ v_1 \quad \dots \quad 1 \quad 1 \\ | \quad \quad \quad | \quad | \\ \dots \quad \quad \quad 1 \quad 1 \\ | \quad \quad \quad | \quad | \\ v_m \quad \quad \quad v_1 \quad 1 \\ | \quad \quad \quad | \quad | \\ \dots \quad \quad \quad \dots \quad 1 \\ | \quad \quad \quad | \quad | \\ v_2 \quad \quad \quad \dots \quad 1 \\ | \quad \quad \quad | \quad | \\ u_1 \quad u_n \quad 1 \end{array}, \quad \frac{X^{(12)}(\underline{u}, \underline{v})}{f(v_{12})a(v_{12})} = \begin{array}{c} \text{---} L \text{---} \\ | \\ \text{---} \Pi \text{---} \\ | \\ \dots \quad \quad \quad 1 \quad 1 \\ | \quad \quad \quad | \quad | \\ \dots \quad \quad \quad 1 \quad 1 \\ | \quad \quad \quad | \quad | \\ v_m \quad \quad \quad v_1 \quad 1 \\ | \quad \quad \quad | \quad | \\ \dots \quad \quad \quad v_2 \quad 1 \\ | \quad \quad \quad | \quad | \\ v_3 \quad \quad \quad \dots \quad 1 \\ | \quad \quad \quad | \quad | \\ u_1 \quad u_n \quad 1 \quad 1 \end{array}$$

■

Note that for the on-shell Bethe ansatz the equations for the unwanted terms look quite similar as (B.7) - and (B.15) (apart from the fact that there are no shifts) and their cancellation is equivalent to the Bethe ansatz equations

$$\chi_i(\underline{u}, \underline{v}) = \frac{a_2(\underline{u}, v_i)}{a_1(\underline{u}, v_i)} \prod_{\substack{k=1 \\ k \neq i}}^m \frac{a(v_{ki})}{a(v_{ik})} = \prod_{k=1}^n \frac{(v_i + \frac{1}{2}) - u_k - \frac{1}{2}}{(v_i + \frac{1}{2}) - u_k + \frac{1}{2}} \prod_{\substack{k=1 \\ k \neq i}}^m \frac{v_{ik} + 1}{v_{ik} - 1} = 1.$$

B.1 The unwanted terms

In our approach of the algebraic Bethe ansatz lemma 2.2 replaces commutation rules of the entries of the monodromy matrix. In the conventional approach one derives them from the Yang-Baxter algebra relations (16) and uses them for the algebraic Bethe ansatz.

The unwanted terms uw^A : We start with (B.3) and use Yang-Baxter relations and lemma 2.2 in the form of (18)

$$\begin{aligned} w^A &= L(\underline{v}) \begin{array}{c} | \quad | \quad | \\ \text{---} \Pi \text{---} \\ | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ u_1 \quad v_i \quad \dots \quad 1 \\ | \quad | \quad | \quad | \\ u'_1 \quad u_n \quad 1 \end{array} = L(\underline{v}) \begin{array}{c} | \quad | \quad | \\ \text{---} \Pi \text{---} \\ | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ u_1 \quad v_i \quad \dots \quad 1 \\ | \quad | \quad | \quad | \\ u'_1 \quad u_n \quad 1 \end{array} \\ &= L(\underline{v}) \begin{array}{c} | \quad | \quad | \\ \text{---} \Pi \text{---} \\ | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ u_1 \quad v_i \quad \dots \quad 1 \\ | \quad | \quad | \quad | \\ u'_1 \quad u_n \quad 1 \end{array} + \sum_{i=1}^m c(u'_1 - v_i) L(\underline{v}) \begin{array}{c} | \quad | \quad | \\ \text{---} \Pi \text{---} \\ | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ \dots \quad \dots \quad 1 \quad 1 \\ | \quad | \quad | \quad | \\ v_i \quad \dots \quad u_1 \quad 1 \\ | \quad | \quad | \quad | \\ u'_1 \quad u_n \quad 1 \end{array} \\ &= \Psi A_Q - uw^A. \end{aligned}$$

Note that the $d(u'_1 - v_i)$ contributions vanish because they produce terms with $\Pi_{\dots\bar{1}} = 0$ (see (30)). This reads in terms of formulas as

$$\begin{aligned} w^A &= (L\Pi)_{\underline{\beta}} \Omega A_Q T_1^{\beta_m}(\underline{u}', v_m) \cdots T_1^{\beta_1}(\underline{u}', v_1) a(u_1 - v_m) \cdots a(u_1 - v_1) \\ &= (L\Pi)_{\underline{\beta}} \Omega (R_{0m}(u'_1 - v_m) \cdots R_{01}(u'_1 - v_1))_{\gamma, \underline{\beta}'}^{\beta, 1} T_1^{\beta_m}(\underline{u}, v_m) \cdots T_1^{\beta_1}(\underline{u}, v_1) (T_Q)_1^\gamma \\ &= (L\Pi)_{\underline{\beta}} \Omega T_1^{\beta_m}(z_m) \cdots T_1^{\beta_1}(z_1) A_Q - uw^A \end{aligned}$$

with

$$\begin{aligned} uw^A &= - \sum_{i=1}^m c(u'_1 - v_i) (L\Pi)_{\underline{\beta}} \Omega (R_{0m}(v_{im}) \cdots \mathbf{P}_{0i} \cdots R_{01}(v_{i1}))_{\gamma, \underline{\beta}'}^{\beta, 1} \\ &\quad \times T_1^{\beta_m}(\underline{u}, v_m) \cdots T_1^{\beta_1}(\underline{u}, v_1) (T_Q)_1^\gamma, \end{aligned}$$

where lemma 2.2 in the form of (18) has been used. We commute the $R_{ik}(v_{ik})$ (for $k < i$) with Π using Yang-Baxter relations and (26) and apply the $\mathring{R}_{ik}(v_{ik})$ to L using (45) such that $\mathring{R}_{ik}(v_{ik}) \rightarrow a(v_{ik})$. Further we use Yang-Baxter relations to the $R_{ik}(v_{ik})$ (for $k > i$) and note that for $R_{0i}(v_{ik})$ only $R_{11}^1(v_{ik}) = a(v_{ik})$ contribute

$$\begin{aligned} uw^A &= - \sum_{i=1}^m c(u'_1 - v_i) (L(v_i, \underline{v}_i) \Pi(v_i, \underline{v}_i))_{\gamma \underline{\beta}_i} \\ &\quad \times \prod_{k=1, k \neq i}^m a(v_{ik}) \Omega A(v_i) \left[T_1^{\beta_m}(v_m) \cdots T_1^{\beta_1}(v_1) \right]_i (T_Q)_1^\gamma \end{aligned} \quad (\text{B.17})$$

$$= (uw_C^A)_{\dot{\gamma}} C_Q^{\dot{\gamma}} + uw_{C_2}^A C_{2,Q}, \quad (\text{B.18})$$

where we use the short notations \underline{v}_i , $\underline{\beta}_i$ and $\left[T_1^{\beta_m}(v_m) \cdots T_1^{\beta_1}(v_1) \right]_i$ which means that v_i , β_i and $T_1^{\beta_i}(v_i)$ are missing, respectively. We can apply (50) $\Omega A(v_i) = a_1(\underline{u}, v_i) \Omega$. The different unwanted terms are due to different values of γ in (B.17). For $\gamma = 1$ the right hand side in (B.17) vanishes because of $\Pi_{\dots\bar{1}} = 0$ (see (28)), for $\gamma = \dot{\gamma} \neq 1, \bar{1}$ this gives uw_C^A and for $\gamma = \bar{1}$ this gives $uw_{C_2}^A$.

First we calculate $(uw_C^A)_{\dot{\gamma}}$ using (29) for $\gamma = \dot{\gamma}$

$$\begin{aligned} (uw_C^A)_{\dot{\gamma}} C_Q^{\dot{\gamma}} &= - \sum_{i=1}^m c(u'_1 - v_i) (L(v_i, \underline{v}_i) \Pi(v_i, \underline{v}_i))_{\dot{\gamma} \underline{\beta}_i} \\ &\quad \times \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik}) a_1(\underline{u}, v_i) \Omega \left[T_1^{\beta_m}(v_m) \cdots T_1^{\beta_1}(v_1) \right]_i (T_Q)_1^{\dot{\gamma}} \\ &= - \sum_{i=1}^m c(u'_1 - v_i) X_{\dot{\gamma}}^{(i)}(\underline{u}, \underline{v}) C_Q^{\dot{\gamma}} \end{aligned}$$

with $X_{\hat{\gamma}}^{(i)}$ defined by (B.11).

$$X_{\hat{\gamma}}^{(i)}(\underline{u}, \underline{v}) = L(v_i, \underline{v}_i)_{\hat{\gamma}\hat{\beta}_i} \Phi_{\hat{\gamma}\hat{\beta}_i}^{\hat{\beta}_i}(\underline{u}, \underline{v}_i) \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik}) a_1(\underline{u}, v_i) \quad (\text{B.19})$$

The remaining unwanted term $uw_{C_2}^A$ comes from $\gamma = \bar{1}$ in (B.17). Using (32) we get

$$\begin{aligned} uw_{C_2}^A C_{2,Q} &= - \sum_{i=1}^m c(u'_1 - v_i) (L(v_i, \underline{v}_i) \Pi(v_i, \underline{v}_i))_{\bar{1}\hat{\beta}_i} \\ &\quad \times \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik}) a_1(\underline{u}, v_i) \Omega \left[T_1^{\beta_m}(v_m) \cdots T_1^{\beta_1}(v_1) \right]_i (T_Q)_1^{\bar{1}} \\ &= - \sum_{i=1}^m c(u'_1 - v_i) \sum_{\substack{j=1 \\ j \neq i}}^m f(v_{ij}) \left(L(v_i, v_j, \underline{v}_{ij}) \hat{C}^{ij} \Pi(\underline{v}_{ij}) \right)_{\hat{\beta}_{ij}} \\ &\quad \times \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik}) a_1(\underline{u}, v_i) \prod_{\substack{k=1 \\ k \neq i, j}}^m a(v_{jk}) a_1(\underline{u}, v_j) \Omega \left[T_1^{\beta_m}(v_m) \cdots T_1^{\beta_1}(v_1) \right]_{ij} C_{2,Q} \\ &= - \sum_{i=1}^m c(u'_1 - v_i) \sum_{\substack{j=1 \\ j \neq i}}^m X^{(ij)}(\underline{v}) C_{2,Q} \end{aligned}$$

with $X^{(ij)}$ defined by (B.16). Therefore we obtain $(uw_{C_2}^{A,i})_{\hat{\gamma}}(\underline{u}, \underline{v})$ and $uw_{C_2}^{A,ij}(\underline{u}, \underline{v})$ in the form of (B.7) and (B.13).

The unwanted terms uw^D : For convenience we add an extra line to Π and consider $\Pi_{\hat{\gamma}\hat{\beta}_i}^{\hat{\beta}_i}(u'_1, \underline{v})$. Using $\Omega T_Q^{\hat{\gamma}} = 0$ (see (51) which also implies that the D -wanted term vanishes), Yang-Baxter and (17) we derive

$$\begin{aligned} 0 &= L(\underline{v}) \quad \begin{array}{c} \hat{\gamma} \\ \Pi \\ \dots \\ u_1 \\ v_i \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} 1 \\ \dots \\ 1 \\ u_1 \\ v_i \\ \dots \\ u_n \end{array} \quad \hat{\gamma} \\ &= L(\underline{v}) \quad \begin{array}{c} \hat{\gamma} \\ \Pi \\ \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} \dots \\ 1 \\ 1 \\ 1 \end{array} \quad \hat{\gamma} \\ &= L(\underline{v}) \quad \begin{array}{c} \hat{\gamma} \\ \Pi \\ \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \hat{\gamma} \\ &\quad + \sum_{i=1}^m c(v_i - u_1) L(\underline{v}) \quad \begin{array}{c} \hat{\gamma} \\ \Pi \\ \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} \dots \\ v_i \\ u_1 \\ \dots \\ u'_1 \end{array} \quad \begin{array}{c} \dots \\ u_n \\ v_i \end{array} \quad \hat{\gamma} \\ &= \Psi(D_Q)_{\hat{\gamma}}^{\hat{\gamma}} - uw^D, \end{aligned}$$

where the dot in the last picture means the the spectral parameter on the left of it is u_1 and on the right of it is v_i . Again the $d(v_i - u_1)$ -terms do not contribute because they produce terms with $\Pi_{\dots\bar{1}} = 0$ (see (30)). In terms of formulas this reads as

$$\begin{aligned} 0 &= L_{\underline{\beta}}(\underline{v}) \left(\Pi(u'_1, \underline{v}) R_{10}(v_1 - u'_1) \cdots R_{m0}(v_m - u'_1) \right)_{\underline{\beta}'\underline{\gamma}'}^{\hat{\gamma}\hat{\beta}} \Omega(T_Q)_{\hat{\gamma}}^{\gamma'} T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \\ &= L_{\underline{\beta}}(\underline{v}) \Pi_{\underline{\gamma}'\underline{\beta}'}^{\hat{\gamma}\hat{\beta}}(u'_1, \underline{v}) \Omega T_{\gamma'_m}^{\beta'_m}(v_m) \cdots T_{\gamma'_1}^{\beta'_1}(v_1) (T_Q)_{\gamma''}^{\gamma'} (R_{10}(v_1 - u_1) \cdots R_{m0}(v_m - u_1))_{1\dots\bar{1},\hat{\gamma}}^{\gamma'',\underline{\gamma}} \\ &= L_{\underline{\beta}}(\underline{v}) \Pi_{\underline{\beta}'}^{\hat{\beta}}(\underline{z}) \Omega T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) (D_Q)_{\hat{\gamma}}^{\hat{\gamma}} - uw^D. \end{aligned}$$

with

$$\begin{aligned} uw^D &= L_{\underline{\beta}}(\underline{v}) \Pi_{\underline{\beta}'}^{\hat{\beta}}(\underline{v}) \Omega T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) (D_Q)_{\hat{\gamma}}^{\hat{\gamma}} \\ &\quad - L_{\underline{\beta}}(\underline{v}) \Pi_{\underline{\gamma}'\underline{\beta}'}^{\hat{\gamma}\hat{\beta}}(u'_1, \underline{v}) \Omega T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) (T_Q)_{\hat{\gamma}}^{\gamma'} - \sum_{i=1}^m c(v_i - u_1) \end{aligned} \quad (\text{B.20})$$

$$\begin{aligned} &\times \Pi_{\underline{\gamma}'\underline{\beta}'}^{\hat{\gamma}\hat{\beta}}(u'_1, \underline{v}) \Omega T_{\gamma'_m}^{\beta'_m}(v_m) \cdots T_{\gamma'_1}^{\beta'_1}(v_1) (T_Q)_{\gamma''}^{\gamma'} (R_{10}(v_{1i}) \cdots \mathbf{P}_{i0} \cdots R_{m0}(v_{mi}))_{1\dots\bar{1},\hat{\gamma}}^{\gamma'',\underline{\gamma}} \\ &\hspace{15em} (\text{B.21}) \end{aligned}$$

$$= (uw_{C_3}^D)_{\hat{\gamma}} C_Q^{\hat{\gamma}} + uw_{C_2}^D C_{2,Q} + (uw_{C_3}^D)_{\hat{\gamma}'} \mathring{\mathbf{C}}^{\hat{\gamma}'\hat{\gamma}'} (C_{3,Q})_{\hat{\gamma}},$$

where lemma 2.2 in the form of (17) has been used. The different unwanted terms are due to different values of γ' in (B.20) and (B.21). For $\gamma' \neq \bar{1}$ in (B.20) the second term cancels the first one. For $\gamma' = \bar{1}$ in (B.20) we get $(uw_{C_3}^D)_{\hat{\gamma}}$

$$(uw_{C_3}^D)_{\hat{\gamma}'} \mathring{\mathbf{C}}^{\hat{\gamma}'\hat{\gamma}'} (C_{3,Q})_{\hat{\gamma}} = -L_{\underline{\beta}}(\underline{v}) \Pi_{\underline{\beta}'}^{\hat{\beta}}(u'_1, \underline{v}) \Omega T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) (T_Q)_{\hat{\gamma}}^{\bar{1}}.$$

Using (34) we get

$$\begin{aligned} &(uw_{C_3}^D)_{\hat{\gamma}'} \mathring{\mathbf{C}}^{\hat{\gamma}'\hat{\gamma}'} \\ &= - \sum_{i=1}^m f(u'_1 - v_i) \left(L(\underline{v}) \mathring{R}_{ii-1} \cdots \mathring{R}_{i1} \mathring{\mathbf{C}}^{0i} \Pi(\underline{v}_i) e_i R_{im} \cdots R_{ii+1} \right)_{\underline{\beta}'}^{\hat{\gamma}} \Omega T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \\ &= - \sum_{i=1}^m f(u'_1 - v_i) (L(v_i, \underline{v}_i) \mathbf{1}_i \Pi(\underline{v}_i))_{\hat{\gamma}'\underline{\beta}'_i} \mathring{\mathbf{C}}^{\hat{\gamma}'\hat{\gamma}'} \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ik}) \Omega A(v_i) \left[T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \right]_i, \end{aligned}$$

where as above we have replaced the R-matrices by $a(v_{ik})$. Finally we obtain with (B.11)

$$(uw_{C_3}^D)_{\hat{\gamma}} = - \sum_{i=1}^m f(u'_1 - v_i) X_{\hat{\gamma}}^{(i)}(\underline{u}, \underline{v}).$$

Therefore we obtain $(uw_{C_3}^{D,i})_{\hat{\gamma}}(\underline{u}, \underline{v})$ in the form of (B.9). The remaining unwanted terms

are due to (B.21)

$$\begin{aligned}
 & (uw_C^D)_{\dot{\gamma}} C_Q^{\dot{\gamma}} + uw_{C_2}^D C_{2,Q} = - \sum_{i=1}^m c(v_i - u_1) \\
 & \quad \times L_{\underline{\beta}}(\underline{v}) \Pi_{\dot{\gamma}'\underline{\beta}'}^{\dot{\gamma}\underline{\beta}}(u'_1, \underline{v}) \Omega T_{\gamma_m}^{\beta'_m}(v_m) \cdots T_{\gamma_1}^{\beta'_1}(v_1) (T_Q)_{\gamma''}^{\gamma'}(R_{10}(v_{1i}) \cdots \mathbf{P}_{i0} \cdots R_{m0}(v_{mi}))_{1 \cdots 1, \dot{\gamma}}^{\gamma'' \gamma} \\
 & = - \sum_{i=1}^m c(v_i - u_1) (L(v'_i, \underline{v}_i) \Pi(u'_1, \underline{v}_i))_{\dot{\gamma}'\underline{\beta}_i} \Omega \left[T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \right]_i \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ki}) a_2(v_i) (T_Q)_1^{\gamma'}
 \end{aligned}$$

where (26), (45), (46) and $\Omega T_{\dot{\gamma}}^{\beta'_i}(v_i) = \delta_{\dot{\gamma}}^{\beta'_i} a_2(v_i) \Omega$ have been used. For $\gamma' = 1$ this vanishes because of (28). For $\gamma' = \dot{\gamma} \neq 1, \bar{1}$ this gives

$$\begin{aligned}
 (uw_C^D)_{\dot{\gamma}} & = - \sum_{i=1}^m c(v_i - u_1) (L(v'_i, \underline{v}_i) \mathbf{1}_i \Pi(\underline{v}_i))_{\dot{\gamma}\underline{\beta}_i} \Omega \left[T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \right]_i \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ki}) a_2(v_i) \\
 & = - \sum_{i=1}^m c(v_i - u_1) X_{\dot{\gamma}}^{(i)}(\underline{u}, \underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v})
 \end{aligned}$$

where $\underline{v}^{(i)}$ means that v_i is replaced by $v'_i = v_i + 2/\nu$ and $\chi_i(\underline{u}, \underline{v})$ is defined by (B.12). For $\gamma' = \bar{1}$ we get using (34) and $c(v_i - u_1) f(u'_1 - v_j) = (c(u_1 - v_i) + f(u'_1 - v_j)) f(v'_{ij})$

$$\begin{aligned}
 uw_{C_2}^D & = - \sum_{i=1}^m c(v_i - u_1) (L(v'_i, \underline{v}_i) \Pi(u'_1, \underline{v}_i))_{\bar{1}\underline{\beta}_i} \Omega \left[T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \right]_i \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ki}) a_2(v_i) \\
 & = - \sum_{i=1}^m c(v_i - u_1) \sum_{\substack{j=1 \\ j \neq i}}^m f(u'_1 - v_j) \left(L(v'_i, v_j, \underline{v}_{ij}) \mathring{\mathbf{C}}^{ij} \Pi(\underline{v}_{ij}) \right)_{\underline{\beta}_i} \\
 & \quad \times \Omega \left[T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \right]_{ij} \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ki}) \prod_{\substack{k=1 \\ k \neq i, j}}^m a(v_{jk}) a_2(v_i) a_1(v_j) \\
 & = - \sum_{i=1}^m \sum_{\substack{j=1 \\ j \neq i}}^m (c(u_1 - v_i) + f(u'_1 - v_j)) X^{(ij)}(\underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v})
 \end{aligned}$$

with $X^{(ij)}$ given by (B.16). Therefore we obtain $(uw_C^{D,i})_{\dot{\gamma}}(\underline{u}, \underline{v})$ and $uw_{C_2}^{D,ij}(\underline{u}, \underline{v})$ in the form of (B.8) and (B.14).

The unwanted terms uw^{A_3} : Using $\Omega T_{Q_1}^{\dot{\gamma}} = 0$ (see (51) which also implies that the A_3 -wanted term vanishes) and Yang-Baxter relations we derive

or in terms of formulas (with $T_{\underline{\beta}, \gamma}^{\gamma', \beta'}(\underline{v}, u) = (R_{10}(v_1 - u) \dots R_{m0}(z_m - u))^{\gamma', \beta'}$ and $T_{\underline{\beta}}^{\beta'}(\underline{u}, v) = (R_{10}(u_1 - v) \dots R_{n0}(u_n - v))^{\beta'}$ where the quantum space indices are suppressed)

$$\begin{aligned}
 0 &= (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}'} T_{\underline{\beta}, \gamma}^{\bar{1}, \beta'}(\underline{v}, u_1) \Omega T_{Q, \bar{1}}^{\gamma}(\underline{u}) T_1^{\beta_m}(\underline{u}', v_m) \dots T_1^{\beta_1}(\underline{u}', v_1) \\
 &= (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta_m}^{\beta_m}(\underline{u}, v_m) \dots T_{\beta_1}^{\beta_1}(\underline{u}, v_1) T_{Q, \gamma}^{\bar{1}}(\underline{u}) T_{1, \dots, 1, \bar{1}}^{\gamma, \beta'}(\underline{v}, u_1) \\
 &= \prod_{k=1}^m (1 + d(v_k - u_1)) \left((L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_1^{\beta_m}(\underline{u}, v_m) \dots T_1^{\beta_1}(\underline{u}, v_1) A_{3, Q}(\underline{u}) - u w^{A_3} \right),
 \end{aligned} \tag{B.22}$$

where the term written down comes from $\gamma = \bar{1}$ in (B.22). It has been used that

$$T_{1, \dots, 1, \bar{1}}^{\bar{1}, \beta'}(\underline{v}, u_1) = (R_{10}(v_1 - u_1) \dots R_{m0}(v_m - u_1))_{1, \dots, 1, \bar{1}}^{\bar{1}, \beta'} = \prod_{k=1}^m (1 + d(v_k - u_1)) \delta_1^{\beta'_k}.$$

The different unwanted terms are due to different values of γ in (B.22): for $\gamma = \hat{\gamma} \neq 1, \bar{1}$ we get $(u w_{C_3}^{A_3})_{\hat{\gamma}'} \mathbf{C}^{\hat{\gamma}'} = (u w_{C_3}^{A_3})^{\hat{\gamma}'}$

$$(u w_{C_3}^{A_3})^{\hat{\gamma}'} \prod_{k=1}^m (1 + d(v_k - u_1)) = - (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta_m}^{\beta_m}(\underline{u}, v_m) \dots T_{\beta_1}^{\beta_1}(\underline{u}, v_1) T_{1, \dots, 1, \bar{1}}^{\hat{\gamma}, \beta'}(\underline{v}, u_1)$$

and for $\gamma = 1$ we get $u w_{C_2}^{A_3}$

$$(u w_{C_2}^{A_3}) \prod_{k=1}^m (1 + d(v_k - u_1)) = - (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta_m}^{\beta_m}(\underline{u}, v_m) \dots T_{\beta_1}^{\beta_1}(\underline{u}, v_1) T_{1, \dots, 1, \bar{1}}^{1, \beta'}(\underline{v}, u_1).$$

The determination of these unwanted terms is not so direct compared to those of the A - and D -unwanted terms. In particular for $u w_{C_2}^{A_3}$ we use more complicated arguments.

To calculate $u w_{C_3}^{A_3}$ we use special components of the Yang Baxter relation (16)

$$R_{\alpha\beta}^{\bar{1}\bar{1}}(1/\nu - 1) T_{\underline{\beta}, \bar{1}}^{\beta\beta}(\underline{v}, u) T_{1, \dots, 1, \hat{\alpha}}^{\alpha\beta'}(\underline{v}, u + 1/\nu - 1) = T_{\underline{\beta}', \alpha}^{1\beta}(\underline{v}, u + 1/\nu - 1) T_{1, \dots, 1, \beta}^{\bar{1}\beta'}(\underline{v}, u) R_{\hat{\alpha}\bar{1}}^{\beta\alpha}(1/\nu - 1).$$

Using $d(1/\nu - 1) = -1$, $T_{1, \dots, 1, \hat{\alpha}}^{\alpha\beta'}(u') = \delta_{\hat{\alpha}}^{\alpha} \mathbf{1}_{1, \dots, 1}^{\beta'}$ for $\alpha \neq 1$ and

$T_{1\dots 1, \beta}^{\bar{1}\beta'}(u) = \delta_{\beta}^{\bar{1}} \mathbf{1}_{1\dots 1}^{\beta'} \prod_{k=1}^m (1 + d(v_k - u))$ we derive

$$\begin{aligned} T_{1\dots 1\bar{1}}^{\hat{\gamma}\beta}(\underline{v}, u) &= -T_{1\dots 1\hat{\gamma}'}^{1\beta}(\underline{v}, u + 1/\nu - 1) \prod_{k=1}^m (1 + d(v_k - u)) \mathbf{C}^{\hat{\gamma}\hat{\gamma}'} \\ &= -\prod_{k=1}^m (1 + d(v_k - u)) \sum_{i=1}^m f(u - v_i) (R_{1a}(v_{1i}) \dots \mathbf{P}_{ia} \dots R_{1a}(v_{mi}))_{1\dots 1\hat{\gamma}'}^{1\beta} \mathbf{C}^{\hat{\gamma}\hat{\gamma}'} \end{aligned}$$

For the last equality (17) and $c(v - (u + 1/\nu - 1)) = f(u - v)$ have been used. Therefore

$$\begin{aligned} (uw_{C_3}^{A_3})_{\hat{\gamma}} &= \sum_{i=1}^m f(u_1 - v_i) (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta'_m}^{\beta m}(\underline{u}, v_m) \dots T_{\beta'_1}^{\beta 1}(\underline{u}, v_1) \\ &\quad \times (R_{1a}(v_{1i}) \dots \mathbf{P}_{ia} \dots R_{1a}(v_{mi}))_{1\dots 1, \hat{\gamma}}^{1\beta} \\ &= \sum_{i=1}^m f(u_1 - v_i) (L(v'_i, \underline{v}_i)\Pi(\underline{v}_i))_{\hat{\gamma}\underline{\beta}_i} \prod_{\substack{k=1 \\ k \neq i}}^m a(v_{ki}) a_2(v_i) \Omega \left[T_1^{\beta m}(v_m) \dots T_1^{\beta 1}(v_1) \right]_i \\ &= \sum_{i=1}^m f(u_1 - v_i) X_{\hat{\gamma}}^{(i)}(\underline{u}, \underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v}) \end{aligned}$$

We obtain $(uw_{C_3}^{A_3, i})_{\hat{\gamma}}(\underline{u}, \underline{v})$ in the form of (B.10). In order to calculate

$$(uw_{C_2}^{A_3}) \prod_{k=1}^m (1 + d(v_k - u_1)) = - (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta'_m}^{\beta m}(\underline{u}, v_m) \dots T_{\beta'_1}^{\beta 1}(\underline{u}, v_1) T_{1\dots 1, \bar{1}}^{1, \beta'}(\underline{v}, u_1) \quad (\text{B.23})$$

we prove

Lemma B.2 *The unwanted term $uw_{C_2}^{A_3}$ is of the form*

$$uw_{C_2}^{A_3} = \sum_{i=1}^m \sum_{j=i, j \neq i}^m (f_{ij}(\underline{v})\Pi(\underline{v}_{ij}))_{\underline{\beta}_{ij}} a_2(v_i) a_2(v_j) \Omega \left[T_1^{\beta m}(v_m) \dots T_1^{\beta 1}(v_1) \right]_{ij}$$

Proof. Using (17) we have

$$\begin{aligned} &\Pi_{\underline{\beta}}^{\hat{\beta}} \Omega T_{\beta'_m}^{\beta m}(\underline{u}, v_m) \dots T_{\beta'_1}^{\beta 1}(\underline{u}, v_1) T_{1\dots 1, \bar{1}}^{1, \beta'}(\underline{v}, u_1) \\ &= \sum_{j=1}^m d(v_j - u_1) \left(\mathring{R}(v_{j-1}) \dots \mathring{R}(v_{j1}) \Pi(v_j, \underline{v}_j) \right)_{\underline{\beta}}^{\hat{\beta}} \Omega \left[T_1^{\beta m}(v_m) \dots T_1^{\beta 1}(v_1) \right]_i T_{\bar{1}}^{\beta j}(v_j) \end{aligned}$$

The $c(v_j - u_1)$ -terms do not contribute because they produce terms like $\Omega B \dots = 0$. The

Yang-Baxter equation for R implies

$$\begin{aligned}
 & \left(\mathring{R}(v_{ji})\Pi(v_{ji}) \right)_{\dots\alpha\beta\dots}^{\mathring{\beta}} \Omega \left[\dots T_{\bar{1}}^{\beta}(v_i)T_{\bar{1}}^{\alpha}(v_j) \dots \right] \\
 &= (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left[\dots T_{\alpha'}^{\alpha}(v_j)T_{\beta'}^{\beta}(v_i) \dots \right] R_{\bar{1}\bar{1}}^{\beta'\alpha'}(v_{ji}) \\
 &= (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left[\dots T_{\bar{1}}^{\alpha}(v_j)T_{\bar{1}}^{\beta}(v_i) \dots \right] (1 + d(v_{ji})) \\
 &+ (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left[\dots T_{\bar{1}}^{\alpha}(v_j)T_{\bar{1}}^{\beta}(v_i) \dots \right] (c(v_{ji}) + d(v_{ji})) \\
 &+ (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left[\dots T_{\bar{\gamma}}^{\alpha}(v_j)T_{\bar{\gamma}}^{\beta}(v_i) \dots \right] d(v_{ji})\mathbf{C}^{\bar{\gamma}\bar{\gamma}} \tag{B.24}
 \end{aligned}$$

Iterating this formula we move the $T_{\bar{1}}$ -operators to the left and finally $\Pi\Omega T_{\bar{1}} \dots = 0$. Therefore $\Pi_{\underline{\beta}}^{\mathring{\beta}}\Omega T_{\bar{1}}^{\beta m}(v_m) \dots T_{\bar{1}}^{\beta_1}(v_1)$ is a sum of terms like

$$\Pi_{\underline{\beta}}^{\mathring{\beta}}\Omega T_{\bar{1}}^{\beta m}(v_m) \dots T_{\bar{\gamma}}^{\beta_i}(v_i)T_{\bar{\gamma}}^{\beta_j}(v_j) \dots T_{\bar{1}}^{\beta_1}(v_1)\mathbf{C}^{\bar{\gamma}\bar{\gamma}}$$

Similar as for $T_{\bar{1}}^{\beta}$ we can move the two $T_{\bar{\gamma}}^{\beta_i}$ -operators to the left using

$$\begin{aligned}
 & \left(\mathring{R}(v_{ji})\Pi(v_{ji}) \right)_{\dots\alpha\beta\dots}^{\mathring{\beta}} \Omega \left(\dots T_{\bar{\gamma}}^{\alpha}(v_i)T_{\bar{1}}^{\beta}(v_j) \dots \right) \\
 &= (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left(\dots T_{\alpha'}^{\alpha}(v_j)T_{\beta'}^{\beta}(v_i) \dots \right) R_{\bar{1}\bar{\gamma}}^{\beta'\alpha'}(v_{ji}) \\
 &= (\Pi(v_{ij}))_{\dots\beta\alpha\dots}^{\mathring{\beta}} \Omega \left(\left(\dots T_{\bar{1}}^{\alpha}(v_j)T_{\bar{\gamma}}^{\beta}(v_i) \dots \right) + \left(\dots T_{\bar{\gamma}}^{\alpha}(v_j)T_{\bar{1}}^{\beta}(v_i) \dots \right) c(v_{ji}) \right)
 \end{aligned}$$

and finally

$$(\Pi)_{\dots\alpha\beta}^{\mathring{\beta}} \Omega T_{\bar{\gamma}}^{\beta}(v_i)T_{\bar{\gamma}}^{\alpha}(v_j) \dots = (\Pi)_{\dots\alpha\beta}^{\mathring{\beta}} \delta_{\bar{\gamma}}^{\beta} \delta_{\bar{\gamma}}^{\alpha} a_2(v_i)a_2(v_j)\Omega \dots$$

■

We calculate $f_{12}(\underline{v})$ and $f_{21}(\underline{v})$, the other $f_{ij}(\underline{v})$ are due to (48) related to $f_{12}(\underline{v})$ by the symmetry

$$(L(v_{ji})\Pi(v_{ji}))_{\dots\beta'\alpha'\dots}^{\mathring{\beta}} \left[\dots T_{\bar{1}}^{\alpha'}(v_i)T_{\bar{1}}^{\beta'}(v_j) \dots \right] = (L(v_{ij})\Pi(v_{ij}))_{\dots\alpha\beta\dots}^{\mathring{\beta}} \left[\dots T_{\bar{1}}^{\beta}(v_j)T_{\bar{1}}^{\alpha}(v_i) \dots \right].$$

We insert in (B.23) the intermediate states $\gamma = 1, 0, \bar{1}$ behind the second R-matrix in the monodromy matrix $T_{1\dots 1, \bar{1}}^{1, \dots}(u, u_1)$

$$\begin{aligned}
 uw_{C_2}^{A_3} \prod_{k=1}^m (1 + d(v_k - u_1)) &= -(L\Pi)_{\underline{\beta}} \Omega T_{\beta'_m}^{\beta m}(\underline{u}, v_m) \dots T_{\beta'_1}^{\beta_1}(\underline{u}, v_1) C_{2Q} \\
 &\times (R(v_1 - u_1)R(v_2 - u_1))_{11, \bar{\gamma}}^{1, \beta'_1 \beta'_2} (R(v_3 - u_1) \dots R(v_m - u_1))_{1\dots 1, \bar{1}}^{\gamma, \beta'_3 \dots \beta'_m}.
 \end{aligned}$$

Similar as in the proof of lemma B.2 one can show that for $\gamma = 1$ there are only contributions to f_{ij} with $2 < i < j$ and for $\gamma = 0$ there are only contributions to f_{1j} or f_{2j} with $2 < j$. So we have to consider only $\gamma = \bar{1}$ where we use

$$(R(v_3 - u_1) \dots R(v_m - u_1))_{1, \dots, \bar{1}}^{\bar{1}, \beta'_3 \dots \beta'_m} = \prod_{k=3}^m (1 + d(v_k - u_1)) \mathbf{1}_{1, \dots, \bar{1}}^{\beta'_3 \dots \beta'_m}$$

to obtain for the contribution of f_{12} and f_{21} to $uw_{C_2}^{A_3} \prod_{k=1}^m (1 + d(v_k - u_1))$

$$\begin{aligned} & - (L\Pi)_{\underline{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_{\beta'_2}^{\beta_2}(v_2) T_{\beta'_1}^{\beta_1}(v_1) C_{2Q} \\ & \quad \times (R(v_1 - u_1) R(v_2 - u_1))_{11, \bar{1}}^{1, \beta'_1 \beta'_2} \prod_{k=3}^m (1 + d(v_k - u_1)) \end{aligned}$$

Using (17) we obtain (similar as in the proof of lemma B.2)

$$\begin{aligned} & \Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_{\beta'_2}^{\beta_2}(v_2) T_{\beta'_1}^{\beta_1}(v_1) (R(v_1 - u_1) R(v_2 - u_1))_{11, \bar{1}}^{1, \beta'_1 \beta'_2} \\ & = c(v_1 - u_1) \left(\dot{R}(v_{21}) \Pi \right)_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_1) T_1^{\beta_1}(v_2) \\ & + c(v_2 - u_1) \Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_2) T_1^{\beta_1}(v_1) a(v_{12}) \\ & + d(v_1 - u_1) \Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_2) T_1^{\beta_1}(v_1) a(v_{12}) \\ & + d(v_2 - u_1) \left(\dot{R}(v_{21}) \Pi \right)_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_1) T_1^{\beta_1}(v_2). \end{aligned}$$

The c -terms do not contribute to f_{12} and f_{21} . For the first d -term we may replace (because of (B.24))

$$\Omega \left[\dots T_1^\alpha(v_2) T_1^\beta(v_1) \right] \rightarrow \Omega \left[\dots T_{\bar{\gamma}}^\alpha(v_2) T_{\bar{\gamma}}^\beta(v_1) \right] f(v_{12}) \mathbf{C}^{\bar{\gamma}\bar{\gamma}} + \dots$$

and obtain

$$\begin{aligned} & d(v_1 - u_1) \Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_2) T_1^{\beta_1}(v_1) a(v_{12}) \\ & = -d(v_1 - u_1) f(v_{12}) \Pi_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_{\bar{\gamma}}^{\beta_2}(v_2) T_{\bar{\gamma}}^{\beta_1}(v_1) a(v_{12}) \mathbf{C}^{\bar{\gamma}\bar{\gamma}} + \dots \end{aligned}$$

where the missing term again does not contribute to f_{12} and f_{21} . Similarly for the second d -term

$$\begin{aligned} & d(v_2 - u_1) \left(\dot{R}(v_{21}) \Pi \right)_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_1) T_1^{\beta_1}(v_2) \\ & = -d(v_2 - u_1) f(v_{21}) \left(\dot{R}(v_{21}) \Pi \right)_{\underline{\beta}}^{\dot{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_{\bar{\gamma}}^{\beta_2}(v_1) T_{\bar{\gamma}}^{\beta_1}(v_2) \mathbf{C}^{\bar{\gamma}\bar{\gamma}} + \dots \end{aligned}$$

In order to move $T_{\tilde{\gamma}}^{\beta}(v_2)T_{\tilde{\gamma}}^{\alpha}(v_1)$ to the left we consider

$$\begin{aligned} & \left(\left(\mathring{R}(v_{32}) \dots \mathring{R}(v_{m2}) \right) \left(\mathring{R}(v_{31}) \dots \mathring{R}(v_{m1}) \right) \Pi \right)_{\underline{\beta}}^{\underline{\beta}} \Omega T_{\tilde{\gamma}}^{\beta_m}(v_2) T_{\tilde{\gamma}}^{\beta_{m-1}}(v_1) T_1^{\beta_2}(v_m) \dots T_1^{\beta_1}(v_3) \mathbf{C}^{\bar{\gamma}\hat{\gamma}} \\ &= \Pi_{\underline{\beta}}^{\underline{\beta}} \Omega T_{\beta'_m}^{\beta_m}(v_m) \dots T_{\beta'_3}^{\beta_3}(v_3) T_{\beta'_2}^{\beta_2}(v_2) T_{\beta'_1}^{\beta_1}(v_1) \\ & \times \left((R(v_{32}) \dots R(v_{m2})) (R(v_{31}) \dots R(v_{m1})) \right)_{1\dots 1, \bar{\gamma}\hat{\gamma}}^{\beta'} \mathbf{C}^{\bar{\gamma}\hat{\gamma}} \\ &= \Pi_{\underline{\beta}}^{\underline{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_{\tilde{\gamma}}^{\beta_2}(v_2) T_{\tilde{\gamma}}^{\beta_1}(v_1) \mathbf{C}^{\bar{\gamma}\hat{\gamma}} + \dots \end{aligned}$$

therefore

$$\begin{aligned} & d(v_1 - u_1) (L\Pi)_{\underline{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_2) T_1^{\beta_1}(v_1) a(v_{12}) \\ &= - \left(L(\underline{v}) \left(\mathring{R}(v_{32}) \dots \mathring{R}(v_{m2}) \right) \left(\mathring{R}(v_{31}) \dots \mathring{R}(v_{m1}) \right) \Pi(\underline{v}_{12}) \mathbf{1}_1 \mathbf{1}_2 \right)_{\underline{\beta}_{12} \beta_1 \beta_2} \\ & \times d(v_1 - u_1) f(v_{12}) a(v_{12}) \Omega T_{\tilde{\gamma}}^{\beta_2}(v_2) T_{\tilde{\gamma}}^{\beta_1}(v_1) T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) \mathbf{C}^{\bar{\gamma}\hat{\gamma}} + \dots \\ &= -d(v_1 - u_1) f(v_{12}) a(v_{12}) \left(L(v'_1, v'_2, \underline{v}_{12}) \mathbf{C}^{12} \Pi(\underline{v}_{12}) \right)_{\underline{\beta}_{12}} \prod_{k=3}^m a(v_{k2}) a(v_{k1}) \\ & \times a_2(v_2) a_2(v_1) \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) + \dots \\ &= -d(v_1 - u_1) X^{(12)}(v'_1, v'_2, \underline{v}_{12}) \frac{a_2(v_2) a_2(v_1) \prod_{k=3}^m a(v_{k2}) a(v_{k1})}{a_1(v'_1) a_1(v'_2) \prod_{k=3}^m a(v'_{2k}) a(v'_{1k})} + \dots \\ &= -d(v_1 - u_1) X^{(12)}(v'_1, v'_2, \underline{v}_{12}) \chi_1(\underline{u}, \underline{v}^{(2)}) \chi_2(\underline{u}, \underline{v}) + \dots \end{aligned}$$

where the missing terms again do not contribute to f_{12} or f_{21} . Similarly

$$\begin{aligned} & d(v_2 - u_1) \left(L\mathring{R}(v_{21}) \Pi \right)_{\underline{\beta}} \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) T_1^{\beta_2}(v_1) T_1^{\beta_1}(v_2) \\ &= - \left(L(\underline{v}) \left(\mathring{R}(v_{21}) \mathring{R}(v_{31}) \dots \mathring{R}(v_{m1}) \right) \left(\mathring{R}(v_{32}) \dots \mathring{R}(v_{m2}) \right) \mathbf{C}^{21} \Pi(\underline{v}_{12}) \right)_{\underline{\beta}_{12}}^{\underline{\beta}} \\ & \times d(v_2 - u_1) f(v_{21}) a_2(v_2) a_2(v_1) \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) + \dots \\ &= -d(v_2 - u_1) f(v_{21}) \left(L(v'_2, v'_1, \underline{v}_{12}) \mathbf{C}^{21} \Pi(\underline{v}_{12}) \right)_{\underline{\beta}_{12}} a(v_{21}) \prod_{k=3}^m a(v_{k2}) a(v_{k1}) \\ & \times a_2(v_2) a_2(v_1) \Omega T_1^{\beta_m}(v_m) \dots T_1^{\beta_3}(v_3) + \dots \\ &= -d(v_2 - u_1) X^{(21)}(v'_2, v'_1, \underline{v}_{12}) \chi_2(\underline{u}, \underline{v}^{(1)}) \chi_1(\underline{u}, \underline{v}) + \dots \end{aligned}$$

Note that $\chi_j(\underline{u}, \underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v}) = \chi_i(\underline{u}, \underline{v}^{(j)}) \chi_j(\underline{u}, \underline{v})$ and

$$X^{(21)}(v'_2, v'_1, \underline{v}_{12}) = -X^{(12)}(v'_1, v'_2, \underline{v}_{12})$$

because of the identities

$$\begin{aligned} L(v'_2, v'_1, \underline{v}_{12}) \mathring{\mathbf{C}}^{21} &= \frac{\mathring{R}_0(v_{21})}{a(v_{21})} L(v'_1, v'_2, \underline{v}_{12}) \mathring{\mathbf{C}}^{12} \\ f(v) \mathring{R}_0(v) &= -a(-v) f(-v). \end{aligned}$$

Finally

$$\begin{aligned}
& uw_{C_2}^{A_3,12} + uw_{C_2}^{A_3,21} \\
&= - \left(\frac{d(v_1 - u_1)}{(1 + d(v_1 - u_1))(1 + d(v_2 - u_1))} - \frac{d(v_2 - u_1)}{(1 + d(v_1 - u_1))(1 + d(v_2 - u_1))} \right) \\
&\times X^{(12)}(v'_1, v'_2, \underline{v}_{12}) \chi_1(\underline{u}, \underline{v}^{(2)}) \chi_2(\underline{u}, \underline{v}) \\
&= (f(u_1 - v_1) - f(u_1 - v_2)) X^{(12)}(v'_1, v'_2, \underline{v}_{12}) \chi_1(\underline{u}, \underline{v}^{(2)}) \chi_2(\underline{u}, \underline{v}) \\
&= (f(u_1 - v_1) X^{(12)}(v'_1, v'_2, \underline{v}_{12}) + f(u_1 - v_2) X^{(21)}(v'_2, v'_1, \underline{v}_{12})) \chi_1(\underline{u}, \underline{v}^{(2)}) \chi_2(\underline{u}, \underline{v})
\end{aligned}$$

we set

$$\begin{aligned}
uw_{C_2}^{A_3,12} &= -f(u_1 - v_2) X^{(12)}(v'_1, v'_2, \underline{v}_{12}) \chi_1(\underline{u}, \underline{v}^{(2)}) \chi_2(\underline{u}, \underline{v}) \\
uw_{C_2}^{A_3,21} &= -f(u_1 - v_1) X^{(21)}(v'_2, v'_1, \underline{v}_{12}) \chi_2(\underline{u}, \underline{v}^{(1)}) \chi_1(\underline{u}, \underline{v}) \\
uw_{C_2}^{A_3,ij} &= -f(u_1 - v_j) X^{(ij)}(v'_i, v'_j, \underline{v}_{ij}) \chi_i(\underline{u}, \underline{v}^{(j)}) \chi_j(\underline{u}, \underline{v})
\end{aligned}$$

which satisfy the desired symmetry. Therefore we obtain $uw_{C_2}^{A_3,ij}(\underline{u}, \underline{v})$ in the form of (B.15).

C Proof of Theorem 4.1

Proof.

1. The weights (74) of the reference state Ω (49) are

$$w = (n = n_0, 0, \dots, 0)$$

In level $k = 1, \dots, [(N - 3) / 2]$ of the Bethe ansatz the weights are changed as

$$w_k \rightarrow w_k - n_k, \quad w_{k+1} \rightarrow w_{k+1} + n_k.$$

This means the states $\Phi_{\underline{\alpha}}^{\underline{\beta}}(\underline{u}, \underline{v})$ of (47) are eigenvectors of the weights. Using in addition (60) for $O(3)$ and (68) for $O(4)$ we obtain $w =$

$$(w_1, \dots, w_{[N/2]}) = \begin{cases} (n - n_1, \dots, n_{[N/2]-1} - n_{[N/2]}, n_{[N/2]}) & \text{for } N \text{ odd} \\ (n - n_1, \dots, n_{[N/2]-2} - n_- - n_+, n_- - n_+) & \text{for } N \text{ even.} \end{cases}$$

2. The proof of the highest weight property

$$\Psi(\underline{v}) M_{\tilde{\gamma}}^1 = \Psi(\underline{v}) M_{\tilde{1}}^{\tilde{\gamma}} = \Psi(\underline{v}) M_{\tilde{1}}^1 = 0$$

uses similar techniques as the derivation of the unwanted terms.

i) We use $\Omega B_{\dot{\gamma}}(v)$, Yang-Baxter relations and apply lemma 2.2 for $v \rightarrow \infty$

$$\begin{aligned}
 0 = & \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} + O(v^{-2}) \\
 & + \sum_{i=1}^m c(v - v_i) \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} + \sum_{i=1}^m c(v_i - v) \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array}
 \end{aligned}$$

The diagrams consist of vertical lines representing particles with various interactions (crossings) and a box labeled Π at the top. Labels include u_1, \dots, u_r at the bottom, v_i and v for crossings, and $\dot{\gamma}$ for the top line.

Multiplied with $L(\underline{v})$ this reads in terms of formulas as

$$\begin{aligned}
 0 = & (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega B_{\dot{\gamma}}(v) T_1^{\beta'_m}(v_m) \cdots T_1^{\beta'_1}(v_1) \\
 = & (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega (R_{0m}(v - v_m) \cdots R_{01}(v - v_1))_{\underline{\gamma}, \underline{\beta}'}^{\underline{\beta}, 1} T_1^{\beta''_m}(v_m) \cdots T_1^{\beta''_1}(v_1) T_{\dot{\gamma}}^{\gamma}(v) \\
 \times & (R_{01}(v_1 - v) \cdots R_{0m}(v_m - v))_{1, \dots, 1, \dot{\gamma}}^{\gamma \underline{\beta}''} + O(v^{-2}).
 \end{aligned}$$

With equations (69), (71) and using similar techniques as for the derivation of uw_C^A and uw_C^D above we obtain

$$0 = \Psi(\underline{v}) M_{\dot{\gamma}}^1 - \sum_{i=1}^m X_{\dot{\gamma}}^{(i)}(\underline{u}, \underline{v}) + \sum_{i=1}^m X_{\dot{\gamma}}^{(i)}(\underline{u}, \underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v})$$

with $X_{\dot{\gamma}}^{(i)}$ and χ_i defined in (B.11) and (B.12). After multiplication with $g(\underline{u}, \underline{v})$ and summation over the \underline{v} the terms cancel each other because of $\chi_i(\underline{u}, \underline{v})g(\underline{u}, \underline{v}) = g(\underline{u}, \underline{v}^{(i)})$.

ii) We consider

$$\begin{aligned}
 & \begin{array}{c} \text{Diagram 1: A box labeled } \Pi \text{ with two input lines from the top. } m \text{ horizontal lines pass through it. The top line is labeled } \gamma \text{ on the left and } \bar{1} \text{ on the right. The } i\text{-th line is labeled } v_i \text{ on the right. The } m\text{-th line is labeled } v \text{ on the right. Below the lines are labels } u_1, \dots, u_r, 1, 1, 1. \end{array} \\
 & = \sum_{i=1}^m d(v_i - v) \begin{array}{c} \text{Diagram 2: Similar to Diagram 1, but the } i\text{-th line is labeled } \bar{1} \text{ on the left and } v \text{ on the right.} \end{array} + O(v^{-2}) \\
 & = \begin{array}{c} \text{Diagram 3: Similar to Diagram 1, but the } i\text{-th line is labeled } \gamma \text{ on the left and } \bar{1} \text{ on the right.} \end{array} + \sum_{i=1}^m d(v_i - v) \begin{array}{c} \text{Diagram 4: Similar to Diagram 1, but the } i\text{-th line is labeled } \gamma \text{ on the left and } \bar{1} \text{ on the right.} \end{array} + O(v^{-2})
 \end{aligned}$$

Multiplied with $L(\underline{v})$ this reads in terms of formulas as

$$\begin{aligned}
 & (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} (R_{10}(v_1 - v) \dots R_{m0}(v_m - v))_{\underline{\beta}, \gamma}^{\hat{\gamma}\underline{\beta}} \Omega T_{\bar{1}}^{\gamma}(v) T_1^{\beta'_m}(v_m) \dots T_1^{\beta'_1}(v_1) \\
 & = (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} (R_{10}(v_1 - v) \dots R_{m0}(v_m - v))_{\bar{1}, \underline{\beta}}^{\hat{\gamma}\underline{\beta}} \Omega T_1^{\beta'_m}(v_m) \dots T_1^{\beta'_1}(v_1) + O(v^{-2}) \\
 & = \Psi(\underline{v}) T_{\bar{1}}^{\hat{\gamma}}(v) + O(v^{-2}) \\
 & + (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} \Omega T_{\beta'_m}^{\beta'_m}(v_m) \dots T_{\beta'_1}^{\beta'_1}(v_1) (R_{10}(v_1 - v) \dots R_{m0}(v_m - v))_{1 \dots 1, \bar{1}}^{\hat{\gamma}\underline{\beta}}.
 \end{aligned}$$

It has been used that only $\gamma = \bar{1}$ contributes because of $\Omega B_2 = \Omega B_3 = 0$. We apply lemma 2.2 for $v \rightarrow \infty$. With equations (69), (71) and using similar techniques as for the derivation of $uw_{C_3}^D$ and $uw_{C_3}^{A_3}$ above we obtain

$$0 = \Psi(\underline{v}) M_{\bar{1}}^{\hat{\gamma}} - \mathbf{C}^{\hat{\gamma}\hat{\gamma}'} \sum_{i=1}^m X_{\hat{\gamma}'}^{(i)}(\underline{u}, \underline{v}) + \mathbf{C}^{\hat{\gamma}\hat{\gamma}'} \sum_{i=1}^m X_{\hat{\gamma}'}^{(i)}(\underline{u}, \underline{v}^{(i)}) \chi_i(\underline{u}, \underline{v}).$$

Again after multiplication with $g(\underline{u}, \underline{v})$ and summation over the \underline{v} the terms cancel each other because of $\chi_i(\underline{u}, \underline{v}) g(\underline{u}, \underline{v}) = g(\underline{u}, \underline{v}^{(i)})$.

iii) We consider

$$0 = \Omega M_{\bar{1}}^1 \dots = \begin{array}{c} \text{Diagram 5: A box labeled } \Pi \text{ with two input lines from the top. } m \text{ horizontal lines pass through it. The } i\text{-th line is labeled } v_i \text{ on the right. The } m\text{-th line is labeled } v \text{ on the right. Below the lines are labels } u_1, \dots, u_r, 1, 1, 1. \end{array} = \begin{array}{c} \text{Diagram 6: Similar to Diagram 5, but the } i\text{-th line is labeled } 1 \text{ on the left and } \bar{1} \text{ on the right.} \end{array}$$

$$\begin{aligned}
 & + \sum_{i=1}^m c(v - v_i) \quad \begin{array}{c} \text{Diagram 1: A box labeled } \Pi \text{ with } m \text{ inputs. The } i\text{-th input is } v_i. \text{ The } i\text{-th output is } \bar{1}. \text{ Other outputs are } 1. \end{array} \\
 & + \sum_{i=1}^m d(v_i - v) \quad \begin{array}{c} \text{Diagram 2: A box labeled } \Pi \text{ with } m \text{ inputs. The } i\text{-th input is } v_i. \text{ The } i\text{-th output is } 1. \text{ Other outputs are } \bar{1}. \end{array} \\
 & + \sum_{i=1}^m c(v_i - v) \quad \begin{array}{c} \text{Diagram 3: A box labeled } \Pi \text{ with } m \text{ inputs. The } i\text{-th input is } v_i. \text{ The } i\text{-th output is } 1. \text{ Other outputs are } \bar{1}. \end{array} \\
 & + O(v^{-2})
 \end{aligned}$$

or in terms of formulas

$$\begin{aligned}
 0 &= \Omega M_{\bar{1}}^1 \dots \\
 &= (L(\underline{v})\Pi(\underline{v}))_{\underline{\beta}} (R(v - v_m) \dots R(v - v_1))_{\underline{\gamma}, \underline{\beta}'}^{\underline{\beta}, 1} \Omega T_{\underline{\beta}''}^{\beta'_m}(v_m) \dots T_{\beta_1'}^{\beta'_1}(v_1) T_{\gamma'}^{\gamma}(v) \\
 &\quad \times (R(v_1 - v) \dots R(v_m - v))_{1, \dots, 1, \bar{1}}^{\gamma', \beta''}.
 \end{aligned}$$

For $v \rightarrow \infty$ we apply lemma 2.2, equations (69) and (71) and obtain

$$0 = \Psi(\underline{v}) M_{\bar{1}}^1 - \sum_{i=1}^m X^{(ij)}(\underline{v}) + \sum_{i=1}^m X^{(ij)}(v'_i, v'_j, \underline{v}_{ij}) \chi_i(\underline{u}, \underline{v}) \chi_j(\underline{u}, \underline{v})$$

where similar techniques as above for the derivation of the unwanted term have been used. Again after multiplication with $g(\underline{u}, \underline{v})$ and summation over the \underline{v} the terms cancel each other because of $\chi_i(\underline{u}, \underline{v}) \chi_j(\underline{u}, \underline{v}) g(\underline{u}, \underline{v}) = g(\underline{u}, \underline{v}^{(ij)})$.

iv) Next we prove

$$\Psi(\underline{v}) M_{\hat{\gamma}'}^{\hat{\gamma}'} = 0, \quad 1 < \hat{\gamma}' < \hat{\gamma} < \bar{1}.$$

We consider

$$\begin{aligned}
 & L_{\underline{\beta}'}(\underline{v}) \Pi_{\underline{\gamma} \underline{\beta}'}^{\hat{\gamma}' \beta'}(v, \underline{v}) \Omega T_1^{\beta_m}(w_m) \dots T_1^{\beta_1}(w_1) T_{\hat{\gamma}'}^{\gamma}(v) + O(v^{-2}) \\
 &= \left(L(\underline{v}) (T^{(1)})_{\hat{\gamma}'}^{\hat{\gamma}'}(v) \right)_{\underline{\beta}'} \Pi_{\underline{\beta}'}^{\beta'}(\underline{v}) \Omega T_{\hat{\gamma}'}^{\gamma}(v) T_1^{\beta_m}(w_m) \dots T_1^{\beta_1}(w_1) + O(v^{-2})
 \end{aligned}$$

where Yang-Baxter rules and (26) have been used. We have also used that by (32) and (17)

$$\begin{aligned}
 \Pi_{\underline{\gamma} \underline{\beta}'}^{\hat{\gamma}' \beta'}(v, \underline{v}) &= \delta_{\hat{\gamma}'}^{\hat{\gamma}'} \Pi_{\underline{\beta}'}^{\beta'}(\underline{v}) + O(v^{-1}) \\
 (R(w_1 - v) \dots R(w_m - v))_{1, \dots, m, 0} &= \mathbf{1}_{1, \dots, m} \mathbf{1}_0 + O(v^{-1}).
 \end{aligned}$$

For $v \rightarrow \infty$ the highest weight condition $L(\underline{v}) (M^{(1)})_{\hat{\gamma}'}^{\hat{\gamma}'} = 0$ implies the claim.

3. The highest weight properties of the weights are obtained as follows. The commutation relation relation (78) reads in the complex basis as

$$[M_\alpha^{\alpha'}, M_\beta^{\beta'}] = -\delta_\alpha^{\beta'} M_\beta^{\alpha'} + \mathbf{C}^{\alpha'\beta'} (\mathbf{C}M)_{\alpha\beta} + M_\alpha^{\beta'} \delta_\beta^{\alpha'} - (\mathbf{M}\mathbf{C})^{\beta'\alpha'} \mathbf{C}_{\alpha\beta}.$$

In particular for $\beta \neq \alpha, \bar{\alpha}$

$$[M_\alpha^\beta, M_\beta^\alpha] = M_\alpha^\alpha - M_\beta^\beta = M_\alpha^\alpha + M_{\bar{\beta}}^\beta.$$

Because of $(M_\alpha^\beta)^\dagger = M_\beta^\alpha$

$$0 \leq M_\alpha^\beta (M_\alpha^\beta)^\dagger = M_\alpha^\beta M_\beta^\alpha = M_\beta^\alpha M_\alpha^\beta + M_\alpha^\alpha - M_\beta^\beta.$$

Applying this to highest weight co-vectors with

$$0 = \Psi M_\beta^\alpha \text{ for } \alpha < \beta$$

we obtain for the weights (74)

$$0 \leq w_\alpha - w_\beta \text{ for } \alpha < \beta \leq N/2.$$

In addition if N is even

$$\begin{aligned} 0 &\leq w_\alpha + w_{\bar{\beta}} \text{ for } \alpha \leq N/2 < \beta \neq \bar{\alpha} \\ &\Rightarrow w_1 \geq w_2 \geq \cdots \geq w_{N/2-1} \geq |w_{N/2}| \end{aligned}$$

and if N is odd

$$\begin{aligned} 0 &\leq w_\alpha \text{ for } \alpha \leq N/2 \text{ because } \Psi M_0^0 = 0 \\ &\Rightarrow w_1 \geq w_2 \geq \cdots \geq w_{N/2} \geq 0. \end{aligned}$$

■

References

- [1] M. Karowski and P. Weisz, Exact form factors in (1+1)-dimensional field theoretic models with soliton behavior, Nucl. Phys. **B139**, 455–476 (1978).
- [2] B. Berg, M. Karowski, and P. Weisz, Construction of Green functions from an exact S matrix, Phys. Rev. **D19**, 2477–2479 (1979).
- [3] F. Smirnov, Form Factors in Completely Integrable Models of Quantum Field Theory, Adv. Series in Math. Phys. **14**, World Scientific (1992).
- [4] H. Babujian, M. Karowski, and A. Zapletal, Matrix Difference Equations and a Nested Bethe ansatz, J. Phys. **A30**, 6425–6450 (1997).

- [5] H. M. Babujian, A. Fring, M. Karowski, and A. Zapletal, Exact form factors in integrable quantum field theories: The sine-Gordon model, *Nucl. Phys.* **B538**, 535–586 (1999).
- [6] H. M. Babujian, A. Foerster, and M. Karowski, The nested $SU(N)$ off-shell Bethe ansatz and exact form factors, *J. Phys.* **A41**, 275202 (2008).
- [7] H. Bethe, On the theory of metals. 1. Eigenvalues and eigenfunctions for the linear atomic chain, *Z. Phys.* **71**, 205–226 (1931).
- [8] L. A. Takhtajan and L. D. Faddeev, The Quantum method of the inverse problem and the Heisenberg XYZ model, *Russ. Math. Surveys* **34**, 11–68 (1979).
- [9] H. M. Babujian, Off-shell Bethe ansatz equation and N point correlators in $SU(2)$ WZNW theory, *J. Phys.* **A26**, 6981–6990 (1993).
- [10] N. Y. Reshetikhin, Jackson-type integrals, bethe vectors, and solutions to a difference analog of the Knizhnik-Zamolodchikov system, *Lett. Math. Phys.* **26**, 153–165 (1992).
- [11] C. N. Yang, Some exact results for the many body problems in one dimension with repulsive delta function interaction, *Phys. Rev. Lett.* **19**, 1312–1314 (1967).
- [12] B. Sutherland, Further Results for the Many-Body Problem in One Dimension, *Phys.Rev.Lett.* **20**, 98–100 (1968).
- [13] B. Sutherland, Model for a multicomponent quantum system, *Phys. Rev. B* **12**(9), 3795–3805 (Nov 1975).
- [14] H. Babujian, M. Karowski, and A. Zapletal, $SU(N)$ Matrix Difference Equations and a Nested Bethe ansatz, hep-th/9611005 (1996).
- [15] V. Tarasov, Algebraic Bethe ansatz for the Izergin-Korepin R Matrix, *Theor. Math. Phys.* **76**, 793–803 (1988).
- [16] M. J. Martins and P. B. Ramos, The algebraic Bethe ansatz for rational braid-monoid lattice models, *Nucl. Phys.* **B500**, 579–620 (1997).
- [17] H. J. de Vega and M. Karowski, Exact Bethe ansatz solution of $O(2N)$ symmetric theories, *Nucl. Phys.* **B280**, 225 (1987).
- [18] H. M. Babujian, A. Foerster, and M. Karowski, Exact form factors of the $O(N)$ Sigma and Gross-Neveu models and the $1/N$ expansion, to be published.
- [19] J. Balog, Form-factors and asymptotic freedom in the $O(3)$ sigma model, *Phys. Lett.* **B300**, 145–151 (1993).
- [20] J. Balog and T. Hauer, Polynomial form-factors in the $O(3)$ nonlinear sigma model, *Phys. Lett.* **B337**, 115–121 (1994).

- [21] L. F. Alday, D. Gaiotto, and Y. Tachikawa, Liouville Correlation Functions from Four-dimensional Gauge Theories, *Lett. Math. Phys.* **91**, 167–197 (2010).
- [22] A. Mironov, A. Morozov, S. Shakirov, and A. Smirnov, Proving AGT conjecture as HS duality: extension to five dimensions, (2011), arXiv:1105.0948 [hep-th].
- [23] I. B. Frenkel and N. Y. Reshetikhin, Quantum affine algebras and holonomic difference equations, *Commun. Math. Phys.* **146**, 1–60 (1992).
- [24] A. B. Zamolodchikov and A. B. Zamolodchikov, Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field models, *Annals Phys.* **120**, 253–291 (1979).
- [25] L. D. Faddeev, E. K. Sklyanin, and L. A. Takhtajan, The quantum inverse problem method. 1, *Theor. Math. Phys.* **40**, 688–706 (1980).
- [26] H. Babujian, A. Foerster, and M. Karowski, Exact form factors in integrable quantum field theories: The scaling $Z(N)$ -Ising model, *Nucl. Phys.* **B736**, 169–198 (2006).
- [27] R. Shankar and E. Witten, The S matrix of the kinks of the $(\bar{\psi}\psi)^2$ model, *Nucl. Phys.* **B141**, 349 (1978).
- [28] M. Karowski and H. J. Thun, Complete S matrix of the $O(2N)$ Gross-Neveu model, *Nucl. Phys.* **B190**, 61–92 (1981).
- [29] H. Babujian, A. Foerster, and M. Karowski, Exact form factors of the $SU(N)$ Gross-Neveu model and $1/N$ expansion, *Nucl. Phys.* **B825**, 396–425 (2010).