

Differenzieren / Kurvendiskussion

Lösung Aufgabe 1:

$$(a) f'(x) = 4(1 - 3x^2)^3 \cdot (-6x) = -24x(1 - 3x^2)^3$$

$$(b) f'(x) = \frac{2x - 5}{2\sqrt{x^2 - 5x + 2}}$$


$$(c) f'(x) = \frac{-\sqrt{x} - (1-x) \cdot \frac{1}{2\sqrt{x}}}{x} = \frac{-2x - (1-x)}{2x^{3/2}} = -\frac{1+x}{2x^{3/2}}$$

$$(d) f'(x) = \frac{1}{2} \sqrt{\frac{x+1}{x-1}} \cdot \frac{(x+1) - (x-1)}{(x+1)^2} = \frac{1}{\sqrt{x-1} \cdot (x+1)^{3/2}}$$

$$(e) f'(x) = -\frac{1}{(\sqrt{x}-1)^2} \cdot \frac{1}{2\sqrt{x}} = -\frac{1}{2\sqrt{x}(\sqrt{x}-1)^2}$$


Lösung Aufgabe 2:

$$(a) \text{ Nullstelle } x_0 = \frac{1}{2}; \text{ y-Achsenabschnitt: } f(0) = 1; f'(x) = -2$$




$$(b) f(x) = \frac{1}{x} \quad \begin{array}{l} f(x) \xrightarrow{x \rightarrow \pm\infty} 0 \\ f(x) \xrightarrow{x \rightarrow \pm 0} \pm \infty \end{array}$$

$$f'(x) = -\frac{1}{x^2} \quad \begin{array}{l} f'(x) \xrightarrow{x \rightarrow \pm\infty} 0 \\ f'(x) \xrightarrow{x \rightarrow \pm 0} -\infty \end{array}$$




(c) $f(x) = \frac{x-1}{x+1}$ $f(x) \xrightarrow{x \rightarrow \pm\infty} 1$
 $f(x) \xrightarrow{x \rightarrow -1 \pm 0} \pm \infty$
 $f(x) = 0$ für $x = 1$
 $f(0) = -1$



$f'(x) = \frac{x+1 - (x-1)}{(x+1)^2} = \frac{2}{(x+1)^2}$ $f'(x) \xrightarrow{x \rightarrow -1 \pm 0} \infty$
 $f'(x) \xrightarrow{x \rightarrow \infty} 0$
 $f(0) = 2$

Lösung Aufgabe 3:

(a) $f'(x) = 1 - \frac{1}{x^2}$ $f'(x) = 0$ für $x = \pm 1$
 $f''(x) = \frac{2}{x^3}$ \rightarrow $x = +1$ ist Minimum
 $x = -1$ ist Maximum
 $f(x) \sim x$ für $x \rightarrow \pm\infty$
 $f(x) \xrightarrow{x \rightarrow \pm 0} \pm \infty$




$$(b) \quad f'(x) = \frac{x^2 + 1 - (x + 2)2x}{(x^2 + 1)^2} = \frac{-x^2 - 4x + 1}{(x^2 + 1)^2}$$

$$\rightarrow f'(x) = 0 \quad \text{wenn} \quad x^2 + 4x - 1 = 0 \Rightarrow x_{1/2} = -2 \pm \sqrt{4 + 1}$$

$$= -2 \pm \sqrt{5}$$

$$f(x) \xrightarrow{x \rightarrow \pm\infty} 0$$


$$f(x) = 0 \quad \text{wenn} \quad x = -2$$



$$(c) \quad f'(x) = \frac{2x}{2\sqrt{x^2 + 2}} = \frac{x}{\sqrt{x^2 + 2}} \Rightarrow f'(x) = 0 \quad \text{für} \quad x = 0$$


$$f(x) \xrightarrow{x \rightarrow \infty} |x|$$

$$f(0) = \sqrt{2}$$




Lösung Aufgabe 4:

$$\begin{aligned}
 \text{(a) } f'(x) &= -\frac{2x}{(x^2 + a^2)^2} \\
 f''(x) &= -\frac{2(x^2 + a^2)^2 - 2x \cdot 2(x^2 + a^2) \cdot 2x}{(x^2 + a^2)^4} \\
 &= -\frac{2(x^2 + a^2) - 8x^2}{(x^2 + a^2)^4} \\
 \Rightarrow f''(x) = 0 &\text{ wenn } 2a^2 - 6x^2 = 0 \Rightarrow x = \pm \frac{a}{\sqrt{3}} \\
 f(x) &\xrightarrow{x \rightarrow \pm\infty} 0 \\
 f(0) &= \frac{1}{a^2}
 \end{aligned}$$



(b) $f(x) = 1 - \frac{1}{x^2 + 1} \rightarrow$ bis auf Verschiebung in y und Vorzeichen wie in (a) mit $a = 1$



Lösung Aufgabe 5:

(a) $f'(x) = 2x \cos(x^2)$

(b) $f'(x) = 2 \sin x \cos x$


(c) $f'(x) = \cos[a \cos(bx)] \cdot (-a) \sin(bx) \cdot b$
 $= -ab \sin bx \cos[a \cos(bx)]$

Lösung Aufgabe 6:



(a) $f(x) \xrightarrow{x \rightarrow +\infty} 0 \quad f(x) \xrightarrow{x \rightarrow -\infty} \infty \quad f(0) = 0$

$$f'(x) = -x^2 e^{-x} + 2x e^{-x} = -x(x-2)e^{-x}$$

$\rightarrow f'(x) = 0$ für $x = 0$ und $x = 2$



(b) $f'(x) = -2ax e^{-ax^2}$




(c)

(Lösung von (b) und (c) nur noch sehr schematisch)

Lösung Aufgabe 7:

$$(a) \quad f(0) = 0; \quad f'(x) = \ln x + 1 \Rightarrow f'(x) = 0 \text{ für } x = \frac{1}{e}$$

$$f''(x) = \frac{1}{x} \neq 0. \quad f(x) = 0 \quad \text{für } x = 0 \text{ und } x = 1$$



$$(b) \quad f(0) = -\infty \quad f(x) = 0 \text{ für } x = (x+1)^2$$

$$\Rightarrow x^2 + x + 1 = 0$$


$$\Rightarrow x = -\frac{1}{2} \pm \sqrt{\frac{1}{4} - 1}$$

→ keine Nullstellen

$$f'(x) = \frac{(x+1)^2}{x} \cdot \frac{(x+1)^2 - x \cdot 2(x+1)}{(x+1)^4} = \frac{x+1-2x}{x(x+1)} = -\frac{x-1}{x(x+1)}$$

$$\Rightarrow f'(x) = 0 \text{ für } x = 1.$$

$$f(x) \sim \ln \frac{1}{x} = -\ln x \text{ für } x \rightarrow \infty$$



Lösung Aufgabe 8:

$$\sinh(\operatorname{arsinh} x) = x$$

$$\Rightarrow \cosh(\operatorname{arsinh} x) \cdot (\operatorname{arsinh} x)' = 1$$

$$\Rightarrow (\operatorname{arsinh} x)' = \frac{1}{\cosh(\operatorname{arsinh} x)}$$

$$= \frac{1}{\sqrt{1 + \sinh^2(\operatorname{arsinh} x)}} = \frac{1}{\sqrt{1 + x^2}}$$



$$\rightarrow (\operatorname{arsinh} x)' = \frac{1}{\sqrt{1 + x^2}}$$

$$\cosh(\operatorname{arcosh} x) = x$$



$$\Rightarrow \sinh(\operatorname{arcosh} x) (\operatorname{arcosh} x)' = 1$$

$$\Rightarrow (\operatorname{arcosh} x)' = \frac{1}{\sinh(\operatorname{arcosh} x)} = \frac{1}{\sqrt{\cosh^2(\operatorname{arcosh} x) - 1}} = \frac{1}{\sqrt{x^2 - 1}}$$

Aufgabe 0a:



Aufgabe 0a:



Aufgabe 0b:

