

Problem Sheet 4
Kraus representation and norms part I

J. Eisert, J. Haferkamp, J. C. Magdalena De La Fuente

1. On the Kraus representation of quantum channels

The operational meaning of Kraus operators can be understood in the following setting in which, for simplicity, we restrict ourselves to quantum channels with the same input and output space $L(\mathcal{X})$. Suppose we apply a unitary U to the joint system and environment in the state $\rho \otimes |0\rangle\langle 0| \in L(\mathcal{X} \otimes \mathcal{Z})$, where $|0\rangle \in \mathcal{Z}$ is some reference state, and then we measure system \mathcal{Z} in the computational basis.

- a) Show that the action of the unitary on the joint system can be written as

$$U(\rho \otimes |0\rangle\langle 0|)U^\dagger = \sum_{kl} E_k \rho E_l^\dagger \otimes |k\rangle\langle l| ,$$

with respect to the basis $\{|i\rangle\}_i$ on the second system.

- b) Now, we perform a von-Neumann measurement on \mathcal{Z} in the same basis. Determine the post-measurement state conditioned on outcome i . What is the probability of obtaining outcome i ?
- c) Give the corresponding operational interpretation of the Kraus operators E_k and the unitary U .
- d) Now, suppose we want to implement a von-Neumann measurement on \mathcal{X} via a global unitary and a von-Neumann measurement on \mathcal{Z} . Characterize the unitaries $U \in U(\mathcal{X} \otimes \mathcal{Z})$ on the joint system that give rise to this situation. Give an example for the case of two qubits.

Finally, we will show some properties of the Kraus representation

- e) Let $\{K_i\}_{i=1}^N$ and $\{\tilde{K}_j\}_{j=1}^N$ be two sets of linear operators in $L(\mathcal{X}, \mathcal{Z})$ fulfilling the completeness relation of Kraus operators. Show that if the two sets are related by a unitary transformations $U \in U(N)$ such that $\tilde{K}_i = \sum_j U_{ij} K_j$, the channels represented by the sets coincide.
- f) Show that all equal-sized Kraus representations of a given channel T are related via a unitary transformation.

Hint: Relate the Kraus representation of two low-rank matrix factorisations of the Choi matrix.

2. ℓ_p -norms

In quantum information we deal with a handful of different matrix spaces such as the set of quantum states and also quantum channels. For quantitative statements we have to equip these spaces with distance measures. Depending on the application and context different distance measures have the desired operational meaning.

A prominent role is played by the so called *Schatten p -norms*. But to set the stage we have to first familiarise ourselves with their analogons on vector spaces, namely *ℓ_p -norms*. For $1 \leq p < \infty$ the ℓ_p -norm on the complex vector space \mathbb{C}^n is defined as

$$\|\bullet\|_{\ell_p} : x \mapsto \|x\|_{\ell_p} := \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} ,$$

and the ℓ_∞ -norm as

$$\|\bullet\|_{\ell_\infty} : x \mapsto \|x\|_{\ell_\infty} := \lim_{p \rightarrow \infty} \|\bullet\|_{\ell_p}.$$

We will now characterise the function $\|\bullet\|_{\ell_p}$ and derive important properties. We begin with an explicit expression for the ℓ_∞ -norm.

a) Show that $\|x\|_{\ell_\infty} = \max_{1 \leq i \leq n} |x_i|$.

For all of what follows the notion of a convex function will be important. Let $D \subset \mathbb{R}$ be a convex set. We say that a function $f : D \rightarrow \mathbb{R}$ is *convex* if

$$f\left(\sum_i a_i x_i\right) \leq \sum_i a_i f(x_i),$$

for all $x_i \in D$ and $a_i \geq 0, i = 1, \dots, m$ such that $\sum_i a_i = 1$.

b) Show that any twice continuously differentiable function on an open interval is convex if and only if its second derivative is everywhere nonnegative.

c) Show that $|\bullet|^p$ is a convex function for $p \geq 1$.

We will now use this fact to show that $\|\bullet\|_{\ell_p}$ is a norm (positive definite, absolutely homogeneous, subadditive aka triangle inequality).

d) Argue that $\|\bullet\|_{\ell_p}$ is positive definite and absolutely homogeneous for $1 \leq p < \infty$ and $p = \infty$.

That was easy. Now comes the hard part; we have to show that the norms satisfy the triangle inequality, i.e.

$$\|x + y\|_{\ell_p} \leq \|x\|_{\ell_p} + \|y\|_{\ell_p}. \tag{1}$$

In fact, the triangle inequality for ℓ_p -norms has even its own name, *Minkowski inequality*. A clever way to prove this inequality is to normalise the right hand side, introduce normalised vectors and then use the convexity of $|\cdot|^p$.

e) Argue that it is sufficient to consider the case $\|x\|_{\ell_p} = \lambda$ and $\|y\|_{\ell_p} = (1 - \lambda)$ with $\lambda \in (0, 1)$ in order to prove the Minkowski inequality.

f) Show the Minkowski inequality for the ℓ_p -norms when $1 \leq p < \infty$.

A crucial property of the ℓ_p -norms is Hölder's inequality. It generalises the Cauchy-Schwarz inequality, which is its special case for $p = 2$. Let $\langle \cdot, \cdot \rangle$ be the Euclidean inner product on \mathbb{C}^n , i.e. $\langle x, y \rangle = \sum_{i=1}^n \bar{x}_i y_i$ with $\bar{\cdot}$ denoting the complex conjugate. Hölder's inequality reads

$$|\langle x, y \rangle| \leq \|x\|_{\ell_p} \|y\|_{\ell_q}, \text{ where } \frac{1}{p} + \frac{1}{q} = 1.$$

Like for the proof of Minkowski's inequality, it will be useful to use normalised vectors in the proof of Hölder's inequality. Furthermore, we will need to first establish the *arithmetic-geometric mean inequality*

$$\prod_{i=1}^n x_i^{a_i} \leq \sum_{i=1}^n a_i x_i \text{ if } x_i \geq 0, a_i \geq 0, \sum_i a_i = 1. \tag{2}$$

g) Show that $-\log$ is a convex function and use this to show the arithmetic-geometric mean inequality, Eq. (2).

h) Now, prove Hölder's inequality for $1 < p < \infty$.

i) Finally, prove Hölder's inequality for $p = 1$.

More generally, for a norm $\|\cdot\|$ on \mathbb{C}^d one can define its dual norm $\|\cdot\|^*$ as

$$\|x\|^* := \sup_{y \in \mathbb{C}^d, \|y\|=1} |\langle x, y \rangle|. \quad (3)$$

j) Show that for every norm $\|\cdot\|$ on \mathbb{C}^d it holds:

$$|\langle x, y \rangle| \leq \|x\| \|y\|^* \quad (4)$$

for all $x, y \in \mathbb{C}^d$.

k) Show that the dual norm $\|\cdot\|_{\ell_p}^*$ of the ℓ_p -norm $\|\cdot\|_{\ell_p}$ is the ℓ_q -norm $\|\cdot\|_{\ell_q}$ with $\frac{1}{p} + \frac{1}{q} = 1$.

Finally, we will show another convenient property of the ℓ_p norms.

l) Show that the ℓ_p norms are ordered in the sense that

$$\|x\|_{\ell_p} \leq \|x\|_{\ell_q}, \text{ for } q \leq p.$$