

Summary Chapter 3 "Statistical Mechanics" 8-3b

microcanonical ensemble : fixed energy U , volume V , and N
 $\hat{=}$ isolated system

partition function $\hat{=}$ volume in phase space (for gas)

$$w(U, V, N) = \int d^3q \int d^3p \quad S(\vec{q}, \vec{p}, t) = \int d^3q \int d^3p \quad 1$$

$H(p, q) = U$ $H(p, q) = U$

$S(\vec{q}, \vec{p}, t) = 1 \hat{=}$ equal a-priori probability
 density of states: $w(U, V, N)$

- entropy $S(U, V, N) = k_B \ln w(U, V, N) \sim N$ extensive
- entropy is maximal at equilibrium $\hat{=}$ 2. law of Th. D.
- $\frac{\partial S(U, V, N)}{\partial U} = \frac{1}{T}$ where T is absolute temperature
- Systems in heat contact have identical $T \hat{=}$ 0. law of Th. D.

canonical ensemble : fixed $T, V, N \hat{=}$ closed system

partition function $Z(T, V, N) = \sum_i e^{-\beta \gamma E_i}$ discrete system

$$= \int d^3q \int d^3p e^{-\beta \gamma E(\vec{q}, \vec{p})}$$

$\beta = 1/k_B T$

$$U = \langle \gamma E \rangle = - \frac{\partial}{\partial \beta} \ln Z = U(T, V, N) \quad \text{caloric equation of state}$$

energy variance $\Delta U^2 = \langle \gamma E^2 \rangle - \langle \gamma E \rangle^2 = \frac{\partial^2 \ln Z}{\partial \beta^2} = k_B T^2 \left. \frac{\partial U}{\partial T} \right|_{V, N}$

$$\left. \frac{\partial U}{\partial T} \right|_{V, N} = C_V \quad \text{heat capacity} \quad C_V \sim N \quad \text{extensive}$$

at const. V

\rightarrow relative energy fluctuations $\frac{\Delta U}{U} \sim \frac{N^{1/2}}{N} \sim N^{-1/2} \rightarrow 0$ as $N \rightarrow \infty$

for $N \rightarrow \infty$ energy fluctuations \rightarrow microcanonical = canonical!

free energy $F(T, V, N) = -k_B T \ln Z(T, V, N)$

at fixed T, V, N the equilibrium state is given by minimal F !

$$S(T, V, N) = - \left. \frac{\partial F(T, V, N)}{\partial T} \right|_V \quad \text{and} \quad \boxed{F = U - TS}$$

total differential of U : $dU = dF + TdS + SdT$ [8-3c]

$$dF(T, V, N) = \left. \frac{\partial F}{\partial T} \right|_{V,N} dT + \left. \frac{\partial F}{\partial V} \right|_{T,N} dV + \left. \frac{\partial F}{\partial N} \right|_{T,V} dN$$

$$\left. \frac{\partial F}{\partial T} \right|_{V,N} = -S(T, V, N), \quad \left. \frac{\partial F}{\partial N} \right|_{T,V} \equiv \mu(T, V, N) \quad (\text{as will see later})$$

$$\rightarrow dU = \left. \frac{\partial F}{\partial V} \right|_{T,N} dV + TdS + \mu dN$$

$$\boxed{dU = -PdV + TdS + \mu dN} \rightarrow \left. \frac{\partial F}{\partial V} \right|_{T,N} = -P(T, V, N)$$

$$\boxed{dU = -\Delta W + \Delta Q + \mu dN}$$

mechanical work heat transfer "chemical work"

1. law of Th.O.

ΔW is not an exact differential, the work W is not a state function

grand-canonical ensemble μ, T, V fixed, N, U fluctuating
 \triangleq open systems

$$Z(\mu, V, T) = \sum_{N=0}^{\infty} e^{\beta N \mu} Z(N, V, T) \quad \text{grand-can. part. function}$$

$$\text{grand potential } \Omega(\mu, V, T) = -k_B T \ln Z(\mu, V, T) = -PV$$

$$\langle N \rangle = \frac{\partial \ln Z(\mu, V, T)}{\beta \partial \mu} = -\frac{\partial \Omega(\mu, V, T)}{\partial \mu}$$

$$\text{particle-number variance } \langle N^2 \rangle - \langle N \rangle^2 = \frac{\partial}{\beta \partial \mu} \frac{\partial}{\beta \partial \mu} \ln Z(\mu, V, T)$$

$$\rightarrow \langle N^2 \rangle - \langle N \rangle^2 = k_B T V \left. \frac{\partial^2 P}{\partial \mu^2} \right|_{T,V} = k_B T N \frac{N}{V} \kappa_T(P, T)$$

$$\rightarrow \frac{\sqrt{\langle N^2 \rangle - \langle N \rangle^2}}{\langle N \rangle} \sim \frac{N^{1/2}}{N} \sim N^{-1/2} \quad \sim N!$$

relative particle number fluctuations vanish for $N \rightarrow \infty$

\rightarrow grand-canonical and canonical ensembles are equivalent!

$$\left. \frac{\partial^2 P}{\partial \mu^2} \right|_{T,V} = \frac{N^2}{V^2} \kappa_T(P, T), \quad \kappa_T = -\frac{1}{V} \left. \frac{\partial V}{\partial P} \right|_{T,N} \geq 0$$

isothermal compressibility

The ideal gas

A) Canonical ensemble

$$Z(N, V, T) = \int d\vec{p}_1 \dots d\vec{p}_N \int d\vec{q}_1 \dots d\vec{q}_N e^{-\beta \gamma E(\vec{q}, \vec{p})} \frac{1}{N! h^{3N}}$$

$$\gamma E = \sum_{i=1}^N \frac{\vec{p}_i^2}{2m}$$

$$\rightarrow Z = \left[\int d^3 p e^{-\vec{p}^2 \beta / 2m} \int d^3 q \right]^N \frac{1}{N! h^{3N}} = \frac{1}{N!} \left(\frac{V}{\lambda_t^3} \right)^N$$

$$\lambda_t = \frac{h}{\sqrt{2\pi m k_B T}} \quad \text{thermal wavelength}$$

$$F(N, V, T) = -k_B T \ln Z(N, V, T) = k_B T N \left[\ln \left(\frac{N \lambda_t^3}{V} \right) - 1 \right]$$

$$\rightarrow S = -\left. \frac{\partial F}{\partial T} \right|_{N, V} = -\frac{F}{T} + \frac{3}{2} k_B N = -k_B N \left[\ln \left(\frac{N \lambda_t^3}{V} \right) - \frac{5}{2} \right]$$

$\rightarrow S(N, V, T) !$

$$P(N, V, T) = -\left. \frac{\partial F}{\partial V} \right|_{N, T} = \frac{k_B T N}{V} \quad \text{ideal equation of state}$$

from $U = F + TS \rightarrow U(N, V, T) = \frac{3}{2} N k_B T$ Caloric Eq. of state

for ideal gas, $U(N, V, T)$ is independent of V !

$$C_V = \frac{\partial U(N, V, T)}{\partial T} = \frac{3}{2} N k_B$$

example of equipartition theorem: per degree of freedom

(= quadratic term in Hamiltonian) $\rightarrow U = \frac{1}{2} k_B T$ and $C_V = \frac{3}{2} N k_B$

B) Microcanonical ensemble: $\omega(N, V, U) = \frac{(V [2mU\pi]^{3/2}/h^3)^N}{N! (3N/2-1)!}$

$$\rightarrow S(N, V, U) = -k_B N \left[\ln \left(\frac{N h^3}{V} \left(\frac{3N}{4mU\pi} \right)^{3/2} \right) - \frac{5}{2} \right]$$

C) Grand canonical ensemble: $Z(\mu, V, T) = \sum_{N=0}^{\infty} e^{\mu N} Z(N, V, T)$

$$Z(\mu, V, T) = \exp \left\{ \frac{V}{\lambda_t^3} e^{\beta \mu} \right\} \rightarrow \Omega(\mu, V, T) = -k_B T \ln Z = -\frac{k_B T V}{\lambda_t^3} e^{\beta \mu}$$

Thermodynamics Foundations

Grand potential explicitly given by $\Omega = -PV$

we also derived $F = \Omega + \mu N = -PV + \mu N$
free energy

we also derived $U = F + TS = TS - PV + \mu N$
internal energy : fundamental equation of thermodynamics

Gibbs free energy $G = F + PV = \mu N$
enthalpy $H = U + PV$

Ω, F, U, G, H are the thermodynamic potentials which are extensive ($\sim N$) and consist of bilinear terms that are products of intensive variables (T, P, μ) and their conjugate extensive state variables (S, V, N).

In principle, other potentials exist, but are not common!
They can all be derived from the fund. eq. for U !

We know that $dU = -PdV + TdS + \mu dN$

so if we have $U(V, S, N)$ in terms of its canonical variables V, S, N we can derive $P(V, S, N)$, $T(V, S, N)$, $\mu(V, S, N)$ and from that all other th.-d. potentials!

as an example: $dF = d(U - TS) = dU - TdS - SdT$
 $dF = -PdV - SdT + \mu dN$

so if we have $F(V, T, N)$ we get $S(V, T, N) = -\frac{\partial F(V, T, N)}{\partial T}$ and similarly $P(V, T, N), \mu(V, T, N) \Rightarrow$ complete information!

canonical forms of th.-d. potentials:

$U(V, S, N), \Omega(V, T, \mu), F(V, T, N), G(P, T, N)$

$H(P, S, N) \dots$ in different ensembles different th.-d. potentials are extremal and therefore useful!